

COMPOSITIONS AND METHODS FOR THE THERAPY AND DIAGNOSIS OF OVARIAN CANCER

CROSS REFERENCE TO RELATED APPLICATIONS

This application is a continuation-in-part of U.S. Application No.
5 09/667,857, filed September 20, 2000, which is a continuation in part of U.S. Application
No. 09/636,801, filed August 10, 2000, which is a continuation-in part of U.S. Application
No. 09/617,747, filed July 17, 2000, which is a continuation-in-part of U.S. Application
No. 09/404,879, filed September 24, 1999, which is a continuation-in-part of U.S.
Application No. 09/338,933, filed June 23, 1999, which is a continuation-in-part of U.S.
10 Application Nos. 09/216,003, filed December 17, 1998, and 09/215,681, filed
December 17, 1998.

TECHNICAL FIELD

The present invention relates generally to ovarian cancer therapy. The
invention is more specifically related to polypeptides comprising at least a portion of an
15 ovarian carcinoma protein, and to polynucleotides encoding such polypeptides, as well as
antibodies and immune system cells that specifically recognize such polypeptides. Such
polypeptides, polynucleotides, antibodies and cells may be used in vaccines and
pharmaceutical compositions for treatment of ovarian cancer.

BACKGROUND OF THE INVENTION

20 Ovarian cancer is a significant health problem for women in the United
States and throughout the world. Although advances have been made in detection and
therapy of this cancer, no vaccine or other universally successful method for prevention or
treatment is currently available. Management of the disease currently relies on a
combination of early diagnosis and aggressive treatment, which may include one or more of
25 a variety of treatments such as surgery, radiotherapy, chemotherapy and hormone therapy.
The course of treatment for a particular cancer is often selected based on a variety of
prognostic parameters, including an analysis of specific tumor markers. However, the use
of established markers often leads to a result that is difficult to interpret, and high mortality
continues to be observed in many cancer patients.

Immunotherapies have the potential to substantially improve cancer treatment and survival. Such therapies may involve the generation or enhancement of an immune response to an ovarian carcinoma antigen. However, to date, relatively few ovarian carcinoma antigens are known and the generation of an immune response against such antigens has not been shown to be therapeutically beneficial.

Accordingly, there is a need in the art for improved methods for identifying ovarian tumor antigens and for using such antigens in the therapy of ovarian cancer. The present invention fulfills these needs and further provides other related advantages.

SUMMARY OF THE INVENTION

Briefly stated, this invention provides compositions and methods for the therapy of cancer, such as ovarian cancer. In one aspect, the present invention provides polypeptides comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished. Within certain embodiments, the ovarian carcinoma protein comprises a sequence that is encoded by a polynucleotide sequence selected from the group consisting of SEQ ID NOs:1-81, 313-331, 359, 366, 379, 385-387, 391 and complements of such polynucleotides.

The present invention further provides polynucleotides that encode a polypeptide as described above or a portion thereof, expression vectors comprising such polynucleotides and host cells transformed or transfected with such expression vectors.

Within other aspects, the present invention provides pharmaceutical compositions and vaccines. Pharmaceutical compositions may comprise a physiologically acceptable carrier or excipient in combination with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ

ID NOs:1-81, 313-331, 359, 366, 379, 385-387 or 391; (ii) a polynucleotide encoding such a polypeptide; (iii) an antibody that specifically binds to such a polypeptide; (iv) an antigen-presenting cell that expresses such a polypeptide and/or (v) a T cell that specifically reacts with such a polypeptide. Vaccines may comprise a non-specific immune response enhancer
 5 in combination with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a
 10 polynucleotide that comprises a sequence recited in any one of SEQ ID NOs:1-81, 313-331, 359, 366, 379, 385-387 or 391; (ii) a polynucleotide encoding such a polypeptide; (iii) an anti-idiotypic antibody that is specifically bound by an antibody that specifically binds to such a polypeptide; (iv) an antigen-presenting cell that expresses such a polypeptide and/or (v) a T cell that specifically reacts with such a polypeptide.

15 The present invention further provides, in other aspects, fusion proteins that comprise at least one polypeptide as described above, as well as polynucleotides encoding such fusion proteins.

Within related aspects, pharmaceutical compositions comprising a fusion protein or polynucleotide encoding a fusion protein in combination with a physiologically
 20 acceptable carrier are provided.

Vaccines are further provided, within other aspects, comprising a fusion protein or polynucleotide encoding a fusion protein in combination with a non-specific immune response enhancer.

Within further aspects, the present invention provides methods for inhibiting
 25 the development of a cancer in a patient, comprising administering to a patient a pharmaceutical composition or vaccine as recited above.

The present invention further provides, within other aspects, methods for stimulating and/or expanding T cells, comprising contacting T cells with (a) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof

that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs:1-387 or 391; (b) a polynucleotide encoding such a polypeptide and/or (c) an antigen presenting cell that expresses such a polypeptide under conditions and for a time sufficient to permit the stimulation and/or expansion of T cells. Such polypeptide, polynucleotide and/or antigen presenting cell(s) may be present within a pharmaceutical composition or vaccine, for use in stimulating and/or expanding T cells in a mammal.

Within other aspects, the present invention provides methods for inhibiting the development of ovarian cancer in a patient, comprising administering to a patient T cells prepared as described above.

Within further aspects, the present invention provides methods for inhibiting the development of ovarian cancer in a patient, comprising the steps of: (a) incubating CD4⁺ and/or CD8⁺ T cells isolated from a patient with one or more of: (i) a polypeptide comprising an immunogenic portion of an ovarian carcinoma protein, or a variant thereof that differs in one or more substitutions, deletions, additions and/or insertions such that the ability of the variant to react with ovarian carcinoma protein-specific antisera is not substantially diminished, wherein the ovarian carcinoma protein comprises an amino acid sequence encoded by a polynucleotide that comprises a sequence recited in any one of SEQ ID NOs: 1-387 or 391; (ii) a polynucleotide encoding such a polypeptide; or (iii) an antigen-presenting cell that expresses such a polypeptide; such that T cells proliferate; and (b) administering to the patient an effective amount of the proliferated T cells, and thereby inhibiting the development of ovarian cancer in the patient. The proliferated cells may be cloned prior to administration to the patient.

The present invention also provides, within other aspects, methods for identifying secreted tumor antigens. Such methods comprise the steps of: (a) implanting tumor cells in an immunodeficient mammal; (b) obtaining serum from the immunodeficient mammal after a time sufficient to permit secretion of tumor antigens into the serum; (c)

immunizing an immunocompetent mammal with the serum; (d) obtaining antiserum from the immunocompetent mammal; and (e) screening a tumor expression library with the antiserum, and therefrom identifying a secreted tumor antigen. A preferred method for identifying a secreted ovarian carcinoma antigen comprises the steps of: (a) implanting
 5 ovarian carcinoma cells in a SCID mouse; (b) obtaining serum from the SCID mouse after a time sufficient to permit secretion of ovarian carcinoma antigens into the serum; (c) immunizing an immunocompetent mouse with the serum; (d) obtaining antiserum from the immunocompetent mouse; and (e) screening an ovarian carcinoma expression library with the antiserum, and therefrom identifying a secreted ovarian carcinoma antigen.

10 The present invention also discloses antibody epitopes recognized by the O8E polyclonal anti-sera which epitopes are presented herein as SEQ ID NOs: 394-415.

Further disclosed by the present invention are 10-mer and 9-mer peptides predicted to bind HLA-0201 which peptides are disclosed herein as SEQ ID NOs: 416-435 and SEQ ID NOs: 436-455, respectively.

15 These and other aspects of the present invention will become apparent upon reference to the following detailed description and attached drawings. All references disclosed herein are hereby incorporated by reference in their entirety as if each was incorporated individually.

BRIEF DESCRIPTION OF THE SEQUENCE IDENTIFIERS AND DRAWINGS

20 SEQ ID NOs:1-71 are ovarian carcinoma antigen polynucleotides shown in Figures 1A-1S.

SEQ ID NOs:72-74 are ovarian carcinoma antigen polynucleotides shown in Figures 2A-2C.

25 SEQ ID NO:75 is the ovarian carcinoma polynucleotide 3g (Figure 4).
 SEQ ID NO:76 is the ovarian carcinoma polynucleotide 3f (Figure 5).
 SEQ ID NO:77 is the ovarian carcinoma polynucleotide 6b (Figure 6).
 SEQ ID NO:78 is the ovarian carcinoma polynucleotide 8e (Figure 7A).
 SEQ ID NO:79 is the ovarian carcinoma polynucleotide 8h (Figure 7B).
 SEQ ID NO:80 is the ovarian carcinoma polynucleotide 12e (Figure 8).

SEQ ID NO:81 is the ovarian carcinoma polynucleotide 12h (Figure 9).

SEQ ID NOs:82-310 are ovarian carcinoma antigen polynucleotides shown in Figures 15A-15EEE.

SEQ ID NO:311 is a full length sequence of ovarian carcinoma
5 polynucleotide O772P.

SEQ ID NO:312 is the O772P amino acid sequence.

SEQ ID NOs:313-384 are ovarian carcinoma antigen polynucleotides.

SEQ ID NOs:385-390 present sequences of O772P forms.

SEQ ID NO:391 is a full length sequence of ovarian carcinoma
10 polynucleotide O8E.

SEQ ID NOs:392-393 are protein sequences encoded by O8E.

SEQ ID NOs:394-415 are peptide sequences corresponding to the OE8
antibody epitopes.

SEQ ID NOs:416-435 are potential HLA-A2 10-mer binding peptides
15 predicted using the full length open-reading frame from OE8.

SEQ ID NOs:436-455 are potential HLA-A2 9-mer binding peptides
predicted using the full length open-reading frame from OE8.

SEQ ID NO:456 is a truncated nucleotide sequence of the full length
Genbank sequence showing homology to O772P

20 SEQ ID NO:457 is the full length Genbank sequence showing significant
homology to O772P

SEQ ID NO:458 is a protein encoding a truncated version of the full length
Genbank sequence showing homology to O772P

25 SEQ ID NO:459 is the full length protein sequence from Genbank showing
significant homology to the protein sequence for O772P

SEQ ID NO:460 encodes a unique N-terminal portion of O772P contained
in residues 1-70.

SEQ ID NO:461 contains unique sequence and encodes residues 1-313 of
SEQ ID NO: 456.

Figures 1A-1S (SEQ ID NOs:1-71) depict partial sequences of polynucleotides encoding representative secreted ovarian carcinoma antigens.

Figures 2A-2C depict full insert sequences for three of the clones of Figure 1. Figure 2A shows the sequence designated O7E (11731; SEQ ID NO:72), Figure 2B shows the sequence designated O9E (11785; SEQ ID NO:73) and Figure 2C shows the sequence designated O8E (13695; SEQ ID NO:74).

Figure 3 presents results of microarray expression analysis of the ovarian carcinoma sequence designated O8E.

Figure 4 presents a partial sequence of a polynucleotide (designated 3g; SEQ ID NO:75) encoding an ovarian carcinoma sequence that is a splice fusion between the human T-cell leukemia virus type I oncoprotein TAX and osteonectin.

Figure 5 presents the ovarian carcinoma polynucleotide designated 3f (SEQ ID NO:76).

Figure 6 presents the ovarian carcinoma polynucleotide designated 6b (SEQ ID NO:77).

Figures 7A and 7B present the ovarian carcinoma polynucleotides designated 8e (SEQ ID NO:78) and 8h (SEQ ID NO:79).

Figure 8 presents the ovarian carcinoma polynucleotide designated 12c (SEQ ID NO:80).

Figure 9 presents the ovarian carcinoma polynucleotide designated 12h (SEQ ID NO:81).

Figure 10 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 3f.

Figure 11 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 6b.

Figure 12 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 8e.

Figure 13 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 12c.

Figure 14 depicts results of microarray expression analysis of the ovarian carcinoma sequence designated 12h.

5 Figures 15A-15EEE depict partial sequences of additional polynucleotides encoding representative secreted ovarian carcinoma antigens (SEQ ID NOs:82-310).

Figure 16 is a diagram illustrating the location of various partial O8E sequences within the full length sequence.

Figure 17 is a graph illustrating the results of epitope mapping studies on
10 O8E protein.

Figure 18 is graph of a fluorescence activated cell sorting (FACS) analysis of O8E cell surface expression.

Figure 19 is graph of a FACS analysis of O8E cell surface expression.

Figure 20 shows FACS analysis results for O8E transfected HEK293 cells
15 demonstrating cell surface expression of O8E.

Figure 21 shows FACS analysis results for SKBR3 breast tumor cells demonstrating cell surface expression of O8E.

Figure 22 shows O8E expression in HEK 293 cells. The cells were probed with anti-O8E rabbit polyclonal antisera #2333L.

20 Figure 23 shows the ELISA analysis of anti-O8E rabbit sera.

Figure 24 shows the ELISA analysis of affinity purified rabbit anti-O8E polyclonal antibody.

Figure 25 is a graph determining antibody internalization of anti-O8E mAb showing that mAbs against amino acids 61-80 induces ligand internalization.

25 DETAILED DESCRIPTION OF THE INVENTION

As noted above, the present invention is generally directed to compositions and methods for the therapy of cancer, such as ovarian cancer. The compositions described herein may include immunogenic polypeptides, polynucleotides encoding such

polypeptides, binding agents such as antibodies that bind to a polypeptide, antigen presenting cells (APCs) and/or immune system cells (*e.g.*, T cells).

Polypeptides of the present invention generally comprise at least an immunogenic portion of an ovarian carcinoma protein or a variant thereof. Certain ovarian carcinoma proteins have been identified using an immunoassay technique, and are referred to herein as ovarian carcinoma antigens. An "ovarian carcinoma antigen" is a protein that is expressed by ovarian tumor cells (preferably human cells) at a level that is at least two fold higher than the level in normal ovarian cells. Certain ovarian carcinoma antigens react detectably (within an immunoassay, such as an ELISA or Western blot) with antisera generated against serum from an immunodeficient animal implanted with a human ovarian tumor. Such ovarian carcinoma antigens are shed or secreted from an ovarian tumor into the sera of the immunodeficient animal. Accordingly, certain ovarian carcinoma antigens provided herein are secreted antigens. Certain nucleic acid sequences of the subject invention generally comprise a DNA or RNA sequence that encodes all or a portion of such a polypeptide, or that is complementary to such a sequence.

The present invention further provides ovarian carcinoma sequences that are identified using techniques to evaluate altered expression within an ovarian tumor. Such sequences may be polynucleotide or protein sequences. Ovarian carcinoma sequences are generally expressed in an ovarian tumor at a level that is at least two fold, and preferably at least five fold, greater than the level of expression in normal ovarian tissue, as determined using a representative assay provided herein. Certain partial ovarian carcinoma polynucleotide sequences are presented herein. Proteins encoded by genes comprising such polynucleotide sequences (or complements thereof) are also considered ovarian carcinoma proteins.

Antibodies are generally immune system proteins, or antigen-binding fragments thereof, that are capable of binding to at least a portion of an ovarian carcinoma polypeptide as described herein. T cells that may be employed within the compositions provided herein are generally T cells (*e.g.*, CD4⁺ and/or CD8⁺) that are specific for such a polypeptide. Certain methods described herein further employ antigen-presenting cells

(such as dendritic cells or macrophages) that express an ovarian carcinoma polypeptide as provided herein.

OVARIAN CARCINOMA POLYNUCLEOTIDES

5 Any polynucleotide that encodes an ovarian carcinoma protein or a portion or other variant thereof as described herein is encompassed by the present invention. Preferred polynucleotides comprise at least 15 consecutive nucleotides, preferably at least 30 consecutive nucleotides, and more preferably at least 45 consecutive nucleotides, that encode a portion of an ovarian carcinoma protein. More preferably, a polynucleotide
10 encodes an immunogenic portion of an ovarian carcinoma protein, such as an ovarian carcinoma antigen. Polynucleotides complementary to any such sequences are also encompassed by the present invention. Polynucleotides may be single-stranded (coding or antisense) or double-stranded, and may be DNA (genomic, cDNA or synthetic) or RNA molecules. Additional coding or non-coding sequences may, but need not, be present
15 within a polynucleotide of the present invention, and a polynucleotide may, but need not, be linked to other molecules and/or support materials.

Polynucleotides may comprise a native sequence (*i.e.*, an endogenous sequence that encodes an ovarian carcinoma protein or a portion thereof) or may comprise a variant of such a sequence. Polynucleotide variants may contain one or more substitutions,
20 additions, deletions and/or insertions such that the immunogenicity of the encoded polypeptide is not diminished, relative to a native ovarian carcinoma protein. The effect on the immunogenicity of the encoded polypeptide may generally be assessed as described herein. Variants preferably exhibit at least about 70% identity, more preferably at least about 80% identity and most preferably at least about 90% identity to a polynucleotide
25 sequence that encodes a native ovarian carcinoma protein or a portion thereof.

The percent identity for two polynucleotide or polypeptide sequences may be readily determined by comparing sequences using computer algorithms well known to those of ordinary skill in the art, such as Megalign, using default parameters. Comparisons between two sequences are typically performed by comparing the sequences over a

comparison window to identify and compare local regions of sequence similarity. A “comparison window” as used herein, refers to a segment of at least about 20 contiguous positions, usually 30 to about 75, or 40 to about 50, in which a sequence may be compared to a reference sequence of the same number of contiguous positions after the two sequences are optimally aligned. Optimal alignment of sequences for comparison may be conducted, for example, using the Megalign program in the Lasergene suite of bioinformatics software (DNASTAR, Inc., Madison, WI), using default parameters. Preferably, the percentage of sequence identity is determined by comparing two optimally aligned sequences over a window of comparison of at least 20 positions, wherein the portion of the polynucleotide or polypeptide sequence in the window may comprise additions or deletions (*i.e.*, gaps) of 20 % or less, usually 5 to 15 %, or 10 to 12%, relative to the reference sequence (which does not contain additions or deletions). The percent identity may be calculated by determining the number of positions at which the identical nucleic acid bases or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the reference sequence (*i.e.*, the window size) and multiplying the results by 100 to yield the percentage of sequence identity.

Variants may also, or alternatively, be substantially homologous to a native gene, or a portion or complement thereof. Such polynucleotide variants are capable of hybridizing under moderately stringent conditions to a naturally occurring DNA sequence encoding a native ovarian carcinoma protein (or a complementary sequence). Suitable moderately stringent conditions include prewashing in a solution of 5 X SSC, 0.5% SDS, 1.0 mM EDTA (pH 8.0); hybridizing at 50°C-65°C, 5 X SSC, overnight; followed by washing twice at 65°C for 20 minutes with each of 2X, 0.5X and 0.2X SSC containing 0.1% SDS.

It will be appreciated by those of ordinary skill in the art that, as a result of the degeneracy of the genetic code, there are many nucleotide sequences that encode a polypeptide as described herein. Some of these polynucleotides bear minimal homology to the nucleotide sequence of any native gene. Nonetheless, polynucleotides that vary due to

differences in codon usage are specifically contemplated by the present invention. Further, alleles of the genes comprising the polynucleotide sequences provided herein are within the scope of the present invention. Alleles are endogenous genes that are altered as a result of one or more mutations, such as deletions, additions and/or substitutions of nucleotides.

- 5 The resulting mRNA and protein may, but need not, have an altered structure or function. Alleles may be identified using standard techniques (such as hybridization, amplification and/or database sequence comparison).

Polynucleotides may be prepared using any of a variety of techniques. For example, an ovarian carcinoma polynucleotide may be identified, as described in more
 10 detail below, by screening a late passage ovarian tumor expression library with antisera generated against sera of immunocompetent mice after injection of such mice with sera from SCID mice implanted with late passage ovarian tumors. Ovarian carcinoma polynucleotides may also be identified using any of a variety of techniques designed to evaluate differential gene expression. Alternatively, polynucleotides may be amplified
 15 from cDNA prepared from ovarian tumor cells. Such polynucleotides may be amplified via polymerase chain reaction (PCR). For this approach, sequence-specific primers may be designed based on the sequences provided herein, and may be purchased or synthesized.

An amplified portion may be used to isolate a full length gene from a suitable library (e.g., an ovarian carcinoma cDNA library) using well known techniques.
 20 Within such techniques, a library (cDNA or genomic) is screened using one or more polynucleotide probes or primers suitable for amplification. Preferably, a library is size-selected to include larger molecules. Random primed libraries may also be preferred for identifying 5' and upstream regions of genes. Genomic libraries are preferred for obtaining introns and extending 5' sequences.

25 For hybridization techniques, a partial sequence may be labeled (e.g., by nick-translation or end-labeling with ^{32}P) using well known techniques. A bacterial or bacteriophage library is then screened by hybridizing filters containing denatured bacterial colonies (or lawns containing phage plaques) with the labeled probe (*see* Sambrook et al., *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratories, Cold Spring

Harbor, NY, 1989). Hybridizing colonies or plaques are selected and expanded, and the DNA is isolated for further analysis. cDNA clones may be analyzed to determine the amount of additional sequence by, for example, PCR using a primer from the partial sequence and a primer from the vector. Restriction maps and partial sequences may be generated to identify one or more overlapping clones. The complete sequence may then be determined using standard techniques, which may involve generating a series of deletion clones. The resulting overlapping sequences are then assembled into a single contiguous sequence. A full length cDNA molecule can be generated by ligating suitable fragments, using well known techniques.

Alternatively, there are numerous amplification techniques for obtaining a full length coding sequence from a partial cDNA sequence. Within such techniques, amplification is generally performed via PCR. Any of a variety of commercially available kits may be used to perform the amplification step. Primers may be designed using, for example, software well known in the art. Primers are preferably 22-30 nucleotides in length, have a GC content of at least 50% and anneal to the target sequence at temperatures of about 68°C to 72°C. The amplified region may be sequenced as described above, and overlapping sequences assembled into a contiguous sequence.

One such amplification technique is inverse PCR (*see* Triglia et al., *Nucl. Acids Res.* 16:8186, 1988), which uses restriction enzymes to generate a fragment in the known region of the gene. The fragment is then circularized by intramolecular ligation and used as a template for PCR with divergent primers derived from the known region. Within an alternative approach, sequences adjacent to a partial sequence may be retrieved by amplification with a primer to a linker sequence and a primer specific to a known region. The amplified sequences are typically subjected to a second round of amplification with the same linker primer and a second primer specific to the known region. A variation on this procedure, which employs two primers that initiate extension in opposite directions from the known sequence, is described in WO 96/38591. Additional techniques include capture PCR (Lagerstrom et al., *PCR Methods Applic.* 1:111-19, 1991) and walking PCR (Parker et

al., *Nucl. Acids. Res.* 19:3055-60, 1991). Other methods employing amplification may also be employed to obtain a full length cDNA sequence.

In certain instances, it is possible to obtain a full length cDNA sequence by analysis of sequences provided in an expressed sequence tag (EST) database, such as that
 5 available from GenBank. Searches for overlapping ESTs may generally be performed using well known programs (*e.g.*, NCBI BLAST searches), and such ESTs may be used to generate a contiguous full length sequence.

Certain nucleic acid sequences of cDNA molecules encoding portions of ovarian carcinoma antigens are provided in Figures 1A-1S (SEQ ID NOS:1 to 71) and
 10 Figures 15A to 15EEE (SEQ ID NOs:82 to 310). The sequences provided in Figures 1A-1S appear to be novel. For sequences in Figures 15A-15EEE, database searches revealed matches having substantial identity. These polynucleotides were isolated by serological screening of an ovarian tumor cDNA expression library, using a technique designed to identify secreted tumor antigens. Briefly, a late passage ovarian tumor expression library
 15 was prepared from a SCID-derived human ovarian tumor (OV9334) in the vector λ -screen (Novagen). The sera used for screening were obtained by injecting immunocompetent mice with sera from SCID mice implanted with one late passage ovarian tumors. This technique permits the identification of cDNA molecules that encode immunogenic portions of secreted tumor antigens.

20 The polynucleotides recited herein, as well as full length polynucleotides comprising such sequences, other portions of such full length polynucleotides, and sequences complementary to all or a portion of such full length molecules, are specifically encompassed by the present invention. It will be apparent to those of ordinary skill in the art that this technique can also be applied to the identification of antigens that are secreted
 25 from other types of tumors.

Other nucleic acid sequences of cDNA molecules encoding portions of ovarian carcinoma proteins are provided in Figures 4-9 (SEQ ID NOs:75-81), as well as SEQ ID NOs:313-384. These sequences were identified by screening a microarray of cDNAs for tumor-associated expression (*i.e.*, expression that is at least five fold greater in

an ovarian tumor than in normal ovarian tissue, as determined using a representative assay provided herein). Such screens were performed using a Synteni microarray (Palo Alto, CA) according to the manufacturer's instructions (and essentially as described by Schena et al., *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997). SEQ ID NOs:311 and 391 provide full length sequences incorporating certain of these nucleic acid sequences.

Any of a variety of well known techniques may be used to evaluate tumor-associated expression of a cDNA. For example, hybridization techniques using labeled polynucleotide probes may be employed. Alternatively, or in addition, amplification techniques such as real-time PCR may be used (see Gibson et al., *Genome Research* 6:995-1001, 1996; Heid et al., *Genome Research* 6:986-994, 1996). Real-time PCR is a technique that evaluates the level of PCR product accumulation during amplification. This technique permits quantitative evaluation of mRNA levels in multiple samples. Briefly, mRNA is extracted from tumor and normal tissue and cDNA is prepared using standard techniques.

Real-time PCR may be performed, for example, using a Perkin Elmer/Applied Biosystems (Foster City, CA) 7700 Prism instrument. Matching primers and fluorescent probes may be designed for genes of interest using, for example, the primer express program provided by Perkin Elmer/Applied Biosystems (Foster City, CA). Optimal concentrations of primers and probes may be initially determined by those of ordinary skill in the art, and control (e.g., β -actin) primers and probes may be obtained commercially from, for example, Perkin Elmer/Applied Biosystems (Foster City, CA). To quantitate the amount of specific RNA in a sample, a standard curve is generated alongside using a plasmid containing the gene of interest. Standard curves may be generated using the Ct values determined in the real-time PCR, which are related to the initial cDNA concentration used in the assay. Standard dilutions ranging from 10^{-10} to 10^{-6} copies of the gene of interest are generally sufficient. In addition, a standard curve is generated for the control sequence. This permits standardization of initial RNA content of a tissue sample to the amount of control for comparison purposes.

Polynucleotide variants may generally be prepared by any method known in the art, including chemical synthesis by, for example, solid phase phosphoramidite chemical synthesis. Modifications in a polynucleotide sequence may also be introduced using standard mutagenesis techniques, such as oligonucleotide-directed site-specific mutagenesis (see Adelman et al., *DNA* 2:183, 1983). Alternatively, RNA molecules may be generated by *in vitro* or *in vivo* transcription of DNA sequences encoding an ovarian carcinoma antigen, or portion thereof, provided that the DNA is incorporated into a vector with a suitable RNA polymerase promoter (such as T7 or SP6). Certain portions may be used to prepare an encoded polypeptide, as described herein. In addition, or alternatively, a portion may be administered to a patient such that the encoded polypeptide is generated *in vivo*.

A portion of a sequence complementary to a coding sequence (*i.e.*, an antisense polynucleotide) may also be used as a probe or to modulate gene expression. cDNA constructs that can be transcribed into antisense RNA may also be introduced into cells or tissues to facilitate the production of antisense RNA. An antisense polynucleotide may be used, as described herein, to inhibit expression of an ovarian carcinoma protein. Antisense technology can be used to control gene expression through triple-helix formation, which compromises the ability of the double helix to open sufficiently for the binding of polymerases, transcription factors or regulatory molecules (see Gee et al., *In* Huber and Carr, *Molecular and Immunologic Approaches*, Futura Publishing Co. (Mt. Kisco, NY; 1994). Alternatively, an antisense molecule may be designed to hybridize with a control region of a gene (*e.g.*, promoter, enhancer or transcription initiation site), and block transcription of the gene; or to block translation by inhibiting binding of a transcript to ribosomes.

Any polynucleotide may be further modified to increase stability *in vivo*. Possible modifications include, but are not limited to, the addition of flanking sequences at the 5' and/or 3' ends; the use of phosphorothioate or 2' O-methyl rather than phosphodiesterase linkages in the backbone; and/or the inclusion of nontraditional bases

such as inosine, queosine and wybutosine, as well as acetyl- methyl-, thio- and other modified forms of adenine, cytidine, guanine, thymine and uridine.

Nucleotide sequences as described herein may be joined to a variety of other nucleotide sequences using established recombinant DNA techniques. For example, a polynucleotide may be cloned into any of a variety of cloning vectors, including plasmids, phagemids, lambda phage derivatives and cosmids. Vectors of particular interest include expression vectors, replication vectors, probe generation vectors and sequencing vectors. In general, a vector will contain an origin of replication functional in at least one organism, convenient restriction endonuclease sites and one or more selectable markers. Other elements will depend upon the desired use, and will be apparent to those of ordinary skill in the art.

Within certain embodiments, polynucleotides may be formulated so as to permit entry into a cell of a mammal, and expression therein. Such formulations are particularly useful for therapeutic purposes, as described below. Those of ordinary skill in the art will appreciate that there are many ways to achieve expression of a polynucleotide in a target cell, and any suitable method may be employed. For example, a polynucleotide may be incorporated into a viral vector such as, but not limited to, adenovirus, adeno-associated virus, retrovirus, or vaccinia or other pox virus (e.g., avian pox virus). Techniques for incorporating DNA into such vectors are well known to those of ordinary skill in the art. A retroviral vector may additionally transfer or incorporate a gene for a selectable marker (to aid in the identification or selection of transduced cells) and/or a targeting moiety, such as a gene that encodes a ligand for a receptor on a specific target cell, to render the vector target specific. Targeting may also be accomplished using an antibody, by methods known to those of ordinary skill in the art.

Other formulations for therapeutic purposes include colloidal dispersion systems, such as macromolecule complexes, nanocapsules, microspheres, beads, and lipid-based systems including oil-in-water emulsions, micelles, mixed micelles, and liposomes. A preferred colloidal system for use as a delivery vehicle *in vitro* and *in vivo* is a liposome

(*i.e.*, an artificial membrane vesicle). The preparation and use of such systems is well known in the art.

OVARIAN CARCINOMA POLYPEPTIDES

5 Within the context of the present invention, polypeptides may comprise at least an immunogenic portion of an ovarian carcinoma protein or a variant thereof, as described herein. As noted above, certain ovarian carcinoma proteins are ovarian carcinoma antigens that are expressed by ovarian tumor cells and react detectably within an immunoassay (such as an ELISA) with antisera generated against serum from an
10 immunodeficient animal implanted with an ovarian tumor. Other ovarian carcinoma proteins are encoded by ovarian carcinoma polynucleotides recited herein. Polypeptides as described herein may be of any length. Additional sequences derived from the native protein and/or heterologous sequences may be present, and such sequences may (but need not) possess further immunogenic or antigenic properties.

15 An "immunogenic portion," as used herein is a portion of an antigen that is recognized (*i.e.*, specifically bound) by a B-cell and/or T-cell surface antigen receptor. Such immunogenic portions generally comprise at least 5 amino acid residues, more preferably at least 10, and still more preferably at least 20 amino acid residues of an ovarian carcinoma protein or a variant thereof. Preferred immunogenic portions are encoded by
20 cDNA molecules isolated as described herein. Further immunogenic portions may generally be identified using well known techniques, such as those summarized in Paul, *Fundamental Immunology*, 3rd ed., 243-247 (Raven Press, 1993) and references cited therein. Such techniques include screening polypeptides for the ability to react with ovarian carcinoma protein-specific antibodies, antisera and/or T-cell lines or clones. As
25 used herein, antisera and antibodies are "ovarian carcinoma protein-specific" if they specifically bind to an ovarian carcinoma protein (*i.e.*, they react with the ovarian carcinoma protein in an ELISA or other immunoassay, and do not react detectably with unrelated proteins). Such antisera, antibodies and T cells may be prepared as described herein, and using well known techniques. An immunogenic portion of a native ovarian

carcinoma protein is a portion that reacts with such antisera, antibodies and/or T-cells at a level that is not substantially less than the reactivity of the full length polypeptide (*e.g.*, in an ELISA and/or T-cell reactivity assay). Such immunogenic portions may react within such assays at a level that is similar to or greater than the reactivity of the full length protein. Such screens may generally be performed using methods well known to those of ordinary skill in the art, such as those described in Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. For example, a polypeptide may be immobilized on a solid support and contacted with patient sera to allow binding of antibodies within the sera to the immobilized polypeptide. Unbound sera may then be removed and bound antibodies detected using, for example, ^{125}I -labeled Protein A.

As noted above, a composition may comprise a variant of a native ovarian carcinoma protein. A polypeptide "variant," as used herein, is a polypeptide that differs from a native ovarian carcinoma protein in one or more substitutions, deletions, additions and/or insertions, such that the immunogenicity of the polypeptide is not substantially diminished. In other words, the ability of a variant to react with ovarian carcinoma protein-specific antisera may be enhanced or unchanged, relative to the native ovarian carcinoma protein, or may be diminished by less than 50%, and preferably less than 20%, relative to the native ovarian carcinoma protein. Such variants may generally be identified by modifying one of the above polypeptide sequences and evaluating the reactivity of the modified polypeptide with ovarian carcinoma protein-specific antibodies or antisera as described herein. Preferred variants include those in which one or more portions, such as an N-terminal leader sequence or transmembrane domain, have been removed. Other preferred variants include variants in which a small portion (*e.g.*, 1-30 amino acids, preferably 5-15 amino acids) has been removed from the N- and/or C-terminal of the mature protein.

Polypeptide variants preferably exhibit at least about 70%, more preferably at least about 90% and most preferably at least about 95% identity to the native polypeptide. Preferably, a variant contains conservative substitutions. A "conservative substitution" is one in which an amino acid is substituted for another amino acid that has

similar properties, such that one skilled in the art of peptide chemistry would expect the secondary structure and hydropathic nature of the polypeptide to be substantially unchanged. Amino acid substitutions may generally be made on the basis of similarity in polarity, charge, solubility, hydrophobicity, hydrophilicity and/or the amphipathic nature of the residues. For example, negatively charged amino acids include aspartic acid and glutamic acid; positively charged amino acids include lysine and arginine; and amino acids with uncharged polar head groups having similar hydrophilicity values include leucine, isoleucine and valine; glycine and alanine; asparagine and glutamine; and serine, threonine, phenylalanine and tyrosine. Other groups of amino acids that may represent conservative changes include: (1) ala, pro, gly, glu, asp, gln, asn, ser, thr; (2) cys, ser, tyr, thr; (3) val, ile, leu, met, ala, phe; (4) lys, arg, his; and (5) phe, tyr, trp, his. A variant may also, or alternatively, contain nonconservative changes. Variants may also (or alternatively) be modified by, for example, the deletion or addition of amino acids that have minimal influence on the immunogenicity, secondary structure and hydropathic nature of the polypeptide.

As noted above, polypeptides may comprise a signal (or leader) sequence at the N-terminal end of the protein which co-translationally or post-translationally directs transfer of the protein. The polypeptide may also be conjugated to a linker or other sequence for ease of synthesis, purification or identification of the polypeptide (*e.g.*, poly-His), or to enhance binding of the polypeptide to a solid support. For example, a polypeptide may be conjugated to an immunoglobulin Fc region.

Polypeptides may be prepared using any of a variety of well known techniques. Recombinant polypeptides encoded by DNA sequences as described above may be readily prepared from the DNA sequences using any of a variety of expression vectors known to those of ordinary skill in the art. Expression may be achieved in any appropriate host cell that has been transformed or transfected with an expression vector containing a DNA molecule that encodes a recombinant polypeptide. Suitable host cells include prokaryotes, yeast and higher eukaryotic cells. Preferably, the host cells employed are *E. coli*, yeast or a mammalian cell line such as COS or CHO. Supernatants from

suitable host/vector systems which secrete recombinant protein or polypeptide into culture media may be first concentrated using a commercially available filter. Following concentration, the concentrate may be applied to a suitable purification matrix such as an affinity matrix or an ion exchange resin. Finally, one or more reverse phase HPLC steps
 5 can be employed to further purify a recombinant polypeptide.

Portions and other variants having fewer than about 100 amino acids, and generally fewer than about 50 amino acids, may also be generated by synthetic means, using techniques well known to those of ordinary skill in the art. For example, such polypeptides may be synthesized using any of the commercially available solid-phase
 10 techniques, such as the Merrifield solid-phase synthesis method, where amino acids are sequentially added to a growing amino acid chain. See Merrifield, *J. Am. Chem. Soc.* 85:2149-2146, 1963. Equipment for automated synthesis of polypeptides is commercially available from suppliers such as Applied BioSystems, Inc. (Foster City, CA), and may be operated according to the manufacturer's instructions.

15 Within certain specific embodiments, a polypeptide may be a fusion protein that comprises multiple polypeptides as described herein, or that comprises one polypeptide as described herein and a known tumor antigen, such as an ovarian carcinoma protein or a variant of such a protein. A fusion partner may, for example, assist in providing T helper epitopes (an immunological fusion partner), preferably T helper epitopes recognized by
 20 humans, or may assist in expressing the protein (an expression enhancer) at higher yields than the native recombinant protein. Certain preferred fusion partners are both immunological and expression enhancing fusion partners. Other fusion partners may be selected so as to increase the solubility of the protein or to enable the protein to be targeted to desired intracellular compartments. Still further fusion partners include affinity tags,
 25 which facilitate purification of the protein.

Fusion proteins may generally be prepared using standard techniques, including chemical conjugation. Preferably, a fusion protein is expressed as a recombinant protein, allowing the production of increased levels, relative to a non-fused protein, in an expression system. Briefly, DNA sequences encoding the polypeptide components may be

assembled separately, and ligated into an appropriate expression vector. The 3' end of the DNA sequence encoding one polypeptide component is ligated, with or without a peptide linker, to the 5' end of a DNA sequence encoding the second polypeptide component so that the reading frames of the sequences are in phase. This permits translation into a single
 5 fusion protein that retains the biological activity of both component polypeptides.

A peptide linker sequence may be employed to separate the first and the second polypeptide components by a distance sufficient to ensure that each polypeptide folds into its secondary and tertiary structures. Such a peptide linker sequence is incorporated into the fusion protein using standard techniques well known in the art.
 10 Suitable peptide linker sequences may be chosen based on the following factors: (1) their ability to adopt a flexible extended conformation; (2) their inability to adopt a secondary structure that could interact with functional epitopes on the first and second polypeptides; and (3) the lack of hydrophobic or charged residues that might react with the polypeptide functional epitopes. Preferred peptide linker sequences contain Gly, Asn and Ser residues.
 15 Other near neutral amino acids, such as Thr and Ala may also be used in the linker sequence. Amino acid sequences which may be usefully employed as linkers include those disclosed in Maratea et al., *Gene* 40:39-46, 1985; Murphy et al., *Proc. Natl. Acad. Sci. USA* 83:8258-8262, 1986; U.S. Patent No. 4,935,233 and U.S. Patent No. 4,751,180. The linker sequence may generally be from 1 to about 50 amino acids in length. Linker sequences are
 20 not required when the first and second polypeptides have non-essential N-terminal amino acid regions that can be used to separate the functional domains and prevent steric interference.

The ligated DNA sequences are operably linked to suitable transcriptional or translational regulatory elements. The regulatory elements responsible for expression of
 25 DNA are located only 5' to the DNA sequence encoding the first polypeptides. Similarly, stop codons required to end translation and transcription termination signals are only present 3' to the DNA sequence encoding the second polypeptide.

Fusion proteins are also provided that comprise a polypeptide of the present invention together with an unrelated immunogenic protein. Preferably the immunogenic

protein is capable of eliciting a recall response. Examples of such proteins include tetanus, tuberculosis and hepatitis proteins (*see*, for example, Stoute et al. *New Engl. J. Med.*, 336:86-91, 1997).

Within preferred embodiments, an immunological fusion partner is derived
 5 from protein D, a surface protein of the gram-negative bacterium *Haemophilus influenza B* (WO 91/18926). Preferably, a protein D derivative comprises approximately the first third of the protein (*e.g.*, the first N-terminal 100-110 amino acids), and a protein D derivative may be lipidated. Within certain preferred embodiments, the first 109 residues of a Lipoprotein D fusion partner is included on the N-terminus to provide the polypeptide with
 10 additional exogenous T-cell epitopes and to increase the expression level in *E. coli* (thus functioning as an expression enhancer). The lipid tail ensures optimal presentation of the antigen to antigen present cells. Other fusion partners include the non-structural protein from influenzae virus, NS1 (hemagglutinin). Typically, the N-terminal 81 amino acids are used, although different fragments that include T-helper epitopes may be used.

15 In another embodiment, the immunological fusion partner is the protein known as LYTA, or a portion thereof (preferably a C-terminal portion). LYTA is derived from *Streptococcus pneumoniae*, which synthesizes an N-acetyl-L-alanine amidase known as amidase LYTA (encoded by the *LytA* gene; *Gene* 43:265-292, 1986). LYTA is an autolysin that specifically degrades certain bonds in the peptidoglycan backbone. The C-
 20 terminal domain of the LYTA protein is responsible for the affinity to the choline or to some choline analogues such as DEAE. This property has been exploited for the development of *E. coli* C-LYTA expressing plasmids useful for expression of fusion proteins. Purification of hybrid proteins containing the C-LYTA fragment at the amino terminus has been described (*see Biotechnology* 10:795-798, 1992). Within a preferred
 25 embodiment, a repeat portion of LYTA may be incorporated into a fusion protein. A repeat portion is found in the C-terminal region starting at residue 178. A particularly preferred repeat portion incorporates residues 188-305.

In general, polypeptides (including fusion proteins) and polynucleotides as described herein are isolated. An "isolated" polypeptide or polynucleotide is one that is

removed from its original environment. For example, a naturally-occurring protein is isolated if it is separated from some or all of the coexisting materials in the natural system. Preferably, such polypeptides are at least about 90% pure, more preferably at least about 95% pure and most preferably at least about 99% pure. A polynucleotide is considered to
 5 be isolated if, for example, it is cloned into a vector that is not a part of the natural environment.

BINDING AGENTS

The present invention further provides agents, such as antibodies and
 10 antigen-binding fragments thereof, that specifically bind to an ovarian carcinoma protein. As used herein, an antibody, or antigen-binding fragment thereof, is said to "specifically bind" to an ovarian carcinoma protein if it reacts at a detectable level (within, for example, an ELISA) with an ovarian carcinoma protein, and does not react detectably with unrelated proteins under similar conditions. As used herein, "binding" refers to a noncovalent
 15 association between two separate molecules such that a "complex" is formed. The ability to bind may be evaluated by, for example, determining a binding constant for the formation of the complex. The binding constant is the value obtained when the concentration of the complex is divided by the product of the component concentrations. In general, two compounds are said to "bind," in the context of the present invention, when the binding
 20 constant for complex formation exceeds about 10^3 L/mol. The binding constant maybe determined using methods well known in the art.

Binding agents may be further capable of differentiating between patients with and without a cancer, such as ovarian cancer, using the representative assays provided herein. In other words, antibodies or other binding agents that bind to a ovarian carcinoma
 25 antigen will generate a signal indicating the presence of a cancer in at least about 20% of patients with the disease, and will generate a negative signal indicating the absence of the disease in at least about 90% of individuals without the cancer. To determine whether a binding agent satisfies this requirement, biological samples (*e.g.*, blood, sera, leukophoresis, urine and/or tumor biopsies) from patients with and without a cancer (as

determined using standard clinical tests) may be assayed as described herein for the presence of polypeptides that bind to the binding agent. It will be apparent that a statistically significant number of samples with and without the disease should be assayed. Each binding agent should satisfy the above criteria; however, those of ordinary skill in the art will recognize that binding agents may be used in combination to improve sensitivity.

Any agent that satisfies the above requirements may be a binding agent. For example, a binding agent may be a ribosome, with or without a peptide component, an RNA molecule or a polypeptide. In a preferred embodiment, a binding agent is an antibody or an antigen-binding fragment thereof. Antibodies may be prepared by any of a variety of techniques known to those of ordinary skill in the art. *See, e.g.,* Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, antibodies can be produced by cell culture techniques, including the generation of monoclonal antibodies as described herein, or via transfection of antibody genes into suitable bacterial or mammalian cell hosts, in order to allow for the production of recombinant antibodies. In one technique, an immunogen comprising the polypeptide is initially injected into any of a wide variety of mammals (*e.g.,* mice, rats, rabbits, sheep or goats). In this step, the polypeptides of this invention may serve as the immunogen without modification. Alternatively, particularly for relatively short polypeptides, a superior immune response may be elicited if the polypeptide is joined to a carrier protein, such as bovine serum albumin or keyhole limpet hemocyanin. The immunogen is injected into the animal host, preferably according to a predetermined schedule incorporating one or more booster immunizations, and the animals are bled periodically. Polyclonal antibodies specific for the polypeptide may then be purified from such antisera by, for example, affinity chromatography using the polypeptide coupled to a suitable solid support.

Monoclonal antibodies specific for an antigenic polypeptide of interest may be prepared, for example, using the technique of Kohler and Milstein, *Eur. J. Immunol.* 6:511-519, 1976, and improvements thereto. Briefly, these methods involve the preparation of immortal cell lines capable of producing antibodies having the desired specificity (*i.e.,* reactivity with the polypeptide of interest). Such cell lines may be

produced, for example, from spleen cells obtained from an animal immunized as described above. The spleen cells are then immortalized by, for example, fusion with a myeloma cell fusion partner, preferably one that is syngeneic with the immunized animal. A variety of fusion techniques may be employed. For example, the spleen cells and myeloma cells may
 5 be combined with a nonionic detergent for a few minutes and then plated at low density on a selective medium that supports the growth of hybrid cells, but not myeloma cells. A preferred selection technique uses HAT (hypoxanthine, aminopterin, thymidine) selection. After a sufficient time, usually about 1 to 2 weeks, colonies of hybrids are observed. Single colonies are selected and their culture supernatants tested for binding activity against the
 10 polypeptide. Hybridomas having high reactivity and specificity are preferred.

Monoclonal antibodies may be isolated from the supernatants of growing hybridoma colonies. In addition, various techniques may be employed to enhance the yield, such as injection of the hybridoma cell line into the peritoneal cavity of a suitable vertebrate host, such as a mouse. Monoclonal antibodies may then be harvested from the
 15 ascites fluid or the blood. Contaminants may be removed from the antibodies by conventional techniques, such as chromatography, gel filtration, precipitation, and extraction. The polypeptides of this invention may be used in the purification process in, for example, an affinity chromatography step.

Within certain embodiments, the use of antigen-binding fragments of
 20 antibodies may be preferred. Such fragments include Fab fragments, which may be prepared using standard techniques. Briefly, immunoglobulins may be purified from rabbit serum by affinity chromatography on Protein A bead columns (Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988) and digested by papain to yield Fab and Fc fragments. The Fab and Fc fragments may be separated by
 25 affinity chromatography on protein A bead columns.

Monoclonal antibodies of the present invention may be coupled to one or more therapeutic agents. Suitable agents in this regard include radionuclides, differentiation inducers, drugs, toxins, and derivatives thereof. Preferred radionuclides include ^{90}Y , ^{123}I , ^{125}I , ^{131}I , ^{186}Re , ^{188}Re , ^{211}At , and ^{212}Bi . Preferred drugs include

methotrexate, and pyrimidine and purine analogs. Preferred differentiation inducers include phorbol esters and butyric acid. Preferred toxins include ricin, abrin, diphtheria toxin, cholera toxin, gclonin, *Pseudomonas* exotoxin, *Shigella* toxin, and pokeweed antiviral protein.

5 A therapeutic agent may be coupled (*e.g.*, covalently bonded) to a suitable monoclonal antibody either directly or indirectly (*e.g.*, via a linker group). A direct reaction between an agent and an antibody is possible when each possesses a substituent capable of reacting with the other. For example, a nucleophilic group, such as an amino or sulfhydryl group, on one may be capable of reacting with a carbonyl-containing group, such as an anhydride or an acid halide, or with an alkyl group containing a good leaving group (*e.g.*, a
10 halide) on the other.

Alternatively, it may be desirable to couple a therapeutic agent and an antibody via a linker group. A linker group can function as a spacer to distance an antibody from an agent in order to avoid interference with binding capabilities. A linker group can
15 also serve to increase the chemical reactivity of a substituent on an agent or an antibody, and thus increase the coupling efficiency. An increase in chemical reactivity may also facilitate the use of agents, or functional groups on agents, which otherwise would not be possible.

It will be evident to those skilled in the art that a variety of bifunctional or
20 polyfunctional reagents, both homo- and hetero-functional (such as those described in the catalog of the Pierce Chemical Co., Rockford, IL), may be employed as the linker group. Coupling may be effected, for example, through amino groups, carboxyl groups, sulfhydryl groups or oxidized carbohydrate residues. There are numerous references describing such methodology, *e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.

25 Where a therapeutic agent is more potent when free from the antibody portion of the immunoconjugates of the present invention, it may be desirable to use a linker group which is cleavable during or upon internalization into a cell. A number of different cleavable linker groups have been described. The mechanisms for the intracellular release of an agent from these linker groups include cleavage by reduction of a

disulfide bond (*e.g.*, U.S. Patent No. 4,489,710, to Spitler), by irradiation of a photolabile bond (*e.g.*, U.S. Patent No. 4,625,014, to Senter et al.), by hydrolysis of derivatized amino acid side chains (*e.g.*, U.S. Patent No. 4,638,045, to Kohn et al.), by serum complement-mediated hydrolysis (*e.g.*, U.S. Patent No. 4,671,958, to Rodwell et al.), and acid-catalyzed
 5 hydrolysis (*e.g.*, U.S. Patent No. 4,569,789, to Blattler et al.).

It may be desirable to couple more than one agent to an antibody. In one embodiment, multiple molecules of an agent are coupled to one antibody molecule. In another embodiment, more than one type of agent may be coupled to one antibody. Regardless of the particular embodiment, immunoconjugates with more than one agent may
 10 be prepared in a variety of ways. For example, more than one agent may be coupled directly to an antibody molecule, or linkers which provide multiple sites for attachment can be used. Alternatively, a carrier can be used.

A carrier may bear the agents in a variety of ways, including covalent bonding either directly or via a linker group. Suitable carriers include proteins such as
 15 albumins (*e.g.*, U.S. Patent No. 4,507,234, to Kato et al.), peptides and polysaccharides such as aminodextran (*e.g.*, U.S. Patent No. 4,699,784, to Shih et al.). A carrier may also bear an agent by noncovalent bonding or by encapsulation, such as within a liposome vesicle (*e.g.*, U.S. Patent Nos. 4,429,008 and 4,873,088). Carriers specific for radionuclide agents include radiohalogenated small molecules and chelating compounds. For example,
 20 U.S. Patent No. 4,735,792 discloses representative radiohalogenated small molecules and their synthesis. A radionuclide chelate may be formed from chelating compounds that include those containing nitrogen and sulfur atoms as the donor atoms for binding the metal, or metal oxide, radionuclide. For example, U.S. Patent No. 4,673,562, to Davison et al. discloses representative chelating compounds and their synthesis.

25 A variety of routes of administration for the antibodies and immunoconjugates may be used. Typically, administration will be intravenous, intramuscular, subcutaneous or in the bed of a resected tumor. It will be evident that the precise dose of the antibody/immunoconjugate will vary depending upon the antibody used, the antigen density on the tumor, and the rate of clearance of the antibody.

Also provided herein are anti-idiotypic antibodies that mimic an immunogenic portion of an ovarian carcinoma protein. Such antibodies may be raised against an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of an ovarian carcinoma protein, using well known techniques. Anti-
 5 idiotypic antibodies that mimic an immunogenic portion of an ovarian carcinoma protein are those antibodies that bind to an antibody, or antigen-binding fragment thereof, that specifically binds to an immunogenic portion of an ovarian carcinoma protein, as described herein.

10 T CELLS

Immunotherapeutic compositions may also, or alternatively, comprise T cells specific for an ovarian carcinoma protein. Such cells may generally be prepared *in vitro* or *ex vivo*, using standard procedures. For example, T cells may be present within (or isolated from) bone marrow, peripheral blood or a fraction of bone marrow or peripheral
 15 blood of a mammal, such as a patient, using a commercially available cell separation system, such as the CEPRATE™ system, available from CellPro Inc., Bothell WA (see also U.S. Patent No. 5,240,856; U.S. Patent No. 5,215,926; WO 89/06280; WO 91/16116 and WO 92/07243). Alternatively, T cells may be derived from related or unrelated humans, non-human animals, cell lines or cultures.

20 T cells may be stimulated with an ovarian carcinoma polypeptide, polynucleotide encoding an ovarian carcinoma polypeptide and/or an antigen presenting cell (APC) that expresses such a polypeptide. Such stimulation is performed under conditions and for a time sufficient to permit the generation of T cells that are specific for the polypeptide. Preferably, an ovarian carcinoma polypeptide or polynucleotide is present
 25 within a delivery vehicle, such as a microsphere, to facilitate the generation of specific T cells.

T cells are considered to be specific for an ovarian carcinoma polypeptide if the T cells kill target cells coated with an ovarian carcinoma polypeptide or expressing a gene encoding such a polypeptide. T cell specificity may be evaluated using any of a

variety of standard techniques. For example, within a chromium release assay or proliferation assay, a stimulation index of more than two fold increase in lysis and/or proliferation, compared to negative controls, indicates T cell specificity. Such assays may be performed, for example, as described in Chen et al., *Cancer Res.* 54:1065-1070, 1994.

- 5 Alternatively, detection of the proliferation of T cells may be accomplished by a variety of known techniques. For example, T cell proliferation can be detected by measuring an increased rate of DNA synthesis (e.g., by pulse-labeling cultures of T cells with tritiated thymidine and measuring the amount of tritiated thymidine incorporated into DNA). Contact with an ovarian carcinoma polypeptide (200 ng/ml - 100 µg/ml, preferably 100
- 10 ng/ml - 25 µg/ml) for 3 - 7 days should result in at least a two fold increase in proliferation of the T cells and/or contact as described above for 2-3 hours should result in activation of the T cells, as measured using standard cytokine assays in which a two fold increase in the level of cytokine release (e.g., TNF or IFN-γ) is indicative of T cell activation (*see* Coligan et al., *Current Protocols in Immunology*, vol. 1, Wiley Interscience (Greene 1998). T cells
- 15 that have been activated in response to an ovarian carcinoma polypeptide, polynucleotide or ovarian carcinoma polypeptide-expressing APC may be CD4⁺ and/or CD8⁺. Ovarian carcinoma polypeptide-specific T cells may be expanded using standard techniques. Within preferred embodiments, the T cells are derived from a patient or a related or unrelated donor and are administered to the patient following stimulation and expansion.

- 20 For therapeutic purposes, CD4⁺ or CD8⁺ T cells that proliferate in response to an ovarian carcinoma polypeptide, polynucleotide or APC can be expanded in number either *in vitro* or *in vivo*. Proliferation of such T cells *in vitro* may be accomplished in a variety of ways. For example, the T cells can be re-exposed to an ovarian carcinoma polypeptide, with or without the addition of T cell growth factors, such as interleukin-2,
- 25 and/or stimulator cells that synthesize an ovarian carcinoma polypeptide. Alternatively, one or more T cells that proliferate in the presence of an ovarian carcinoma polypeptide can be expanded in number by cloning. Methods for cloning cells are well known in the art, and include limiting dilution. Following expansion, the cells may be administered back to the patient as described, for example, by Chang et al., *Crit. Rev. Oncol. Hematol.* 22:213,

1996.

PHARMACEUTICAL COMPOSITIONS AND VACCINES

Within certain aspects, polypeptides, polynucleotides, binding agents and/or immune system cells as described herein may be incorporated into pharmaceutical compositions or vaccines. Pharmaceutical compositions comprise one or more such compounds or cells and a physiologically acceptable carrier. Vaccines may comprise one or more such compounds or cells and a non-specific immune response enhancer. A non-specific immune response enhancer may be any substance that enhances an immune response to an exogenous antigen. Examples of non-specific immune response enhancers include adjuvants, biodegradable microspheres (*e.g.*, polylactic galactide) and liposomes (into which the compound is incorporated; *see e.g.*, Fullerton, U.S. Patent No. 4,235,877). Vaccine preparation is generally described in, for example, M.F. Powell and M.J. Newman, eds., "Vaccine Design (the subunit and adjuvant approach)," Plenum Press (NY, 1995). Pharmaceutical compositions and vaccines within the scope of the present invention may also contain other compounds, which may be biologically active or inactive. For example, one or more immunogenic portions of other tumor antigens may be present, either incorporated into a fusion polypeptide or as a separate compound within the composition or vaccine.

A pharmaceutical composition or vaccine may contain DNA encoding one or more of the polypeptides as described above, such that the polypeptide is generated *in situ*. As noted above, the DNA may be present within any of a variety of delivery systems known to those of ordinary skill in the art, including nucleic acid expression systems, bacteria and viral expression systems. Appropriate nucleic acid expression systems contain the necessary DNA sequences for expression in the patient (such as a suitable promoter and terminating signal). Bacterial delivery systems involve the administration of a bacterium (such as *Bacillus-Calmette-Guerrin*) that expresses an immunogenic portion of the polypeptide on its cell surface. In a preferred embodiment, the DNA may be introduced using a viral expression system (*e.g.*, vaccinia or other pox virus, retrovirus, or adenovirus),

which may involve the use of a non-pathogenic (defective), replication competent virus. Suitable systems are disclosed, for example, in Fisher-Hoch et al., *PNAS* 86:317-321, 1989; Flexner et al., *Ann. N Y Acad. Sci.* 569:86-103, 1989; Flexner et al., *Vaccine* 8:17-21, 1990; U.S. Patent Nos. 4,603,112, 4,769,330, and 5,017,487; WO 89/01973; U.S. Patent
 5 No. 4,777,127; GB 2,200,651; EP 0,345,242; WO 91/02805; Berkner, *Biotechniques* 6:616-627, 1988; Rosenfeld et al., *Science* 252:431-434, 1991; Kolls et al., *PNAS* 91:215-219, 1994; Kass-Eisler et al., *PNAS* 90:11498-11502, 1993; Guzman et al., *Circulation* 88:2838-2848, 1993; and Guzman et al., *Cir. Res.* 73:1202-1207, 1993. Techniques for incorporating DNA into such expression systems are well known to those of
 10 ordinary skill in the art. The DNA may also be "naked," as described, for example, in Ulmer et al., *Science* 259:1745-1749, 1993 and reviewed by Cohen, *Science* 259:1691-1692, 1993. The uptake of naked DNA may be increased by coating the DNA onto biodegradable beads, which are efficiently transported into the cells.

While any suitable carrier known to those of ordinary skill in the art may be
 15 employed in the pharmaceutical compositions of this invention, the type of carrier will vary depending on the mode of administration. Compositions of the present invention may be formulated for any appropriate manner of administration, including for example, topical, oral, nasal, intravenous, intracranial, intraperitoneal, subcutaneous or intramuscular administration. For parenteral administration, such as subcutaneous injection, the carrier
 20 preferably comprises water, saline, alcohol, a fat, a wax or a buffer. For oral administration, any of the above carriers or a solid carrier, such as mannitol, lactose, starch, magnesium stearate, sodium saccharine, talcum, cellulose, glucose, sucrose, and magnesium carbonate, may be employed. Biodegradable microspheres (*e.g.*, polylactate polyglycolate) may also be employed as carriers for the pharmaceutical compositions of
 25 this invention. Suitable biodegradable microspheres are disclosed, for example, in U.S. Patent Nos. 4,897,268 and 5,075,109.

Such compositions may also comprise buffers (*e.g.*, neutral buffered saline or phosphate buffered saline), carbohydrates (*e.g.*, glucose, mannose, sucrose or dextrans), mannitol, proteins, polypeptides or amino acids such as glycine, antioxidants, chelating

agents such as EDTA or glutathione, adjuvants (*e.g.*, aluminum hydroxide) and/or preservatives. Alternatively, compositions of the present invention may be formulated as a lyophilizate. Compounds may also be encapsulated within liposomes using well known technology.

5 Any of a variety of non-specific immune response enhancers may be employed in the vaccines of this invention. For example, an adjuvant may be included. Most adjuvants contain a substance designed to protect the antigen from rapid catabolism, such as aluminum hydroxide or mineral oil, and a stimulator of immune responses, such as lipid A, *Bordetella pertussis* or *Mycobacterium tuberculosis* derived proteins. Suitable
 10 adjuvants are commercially available as, for example, Freund's Incomplete Adjuvant and Complete Adjuvant (Difco Laboratories, Detroit, MI), Merck Adjuvant 65 (Merck and Company, Inc., Rahway, NJ), alum, biodegradable microspheres, monophosphoryl lipid A and quil A. Cytokines, such as GM-CSF or interleukin-2, -7, or -12, may also be used as adjuvants.

15 Within the vaccines provided herein, the adjuvant composition is preferably designed to induce an immune response predominantly of the Th1 type. High levels of Th1-type cytokines (*e.g.*, IFN- γ , IL-2 and IL-12) tend to favor the induction of cell mediated immune responses to an administered antigen. In contrast, high levels of Th2-type cytokines (*e.g.*, IL-4, IL-5, IL-6, IL-10 and TNF- β) tend to favor the induction of
 20 humoral immune responses. Following application of a vaccine as provided herein, a patient will support an immune response that includes Th1- and Th2-type responses. Within a preferred embodiment, in which a response is predominantly Th1-type, the level of Th1-type cytokines will increase to a greater extent than the level of Th2-type cytokines. The levels of these cytokines may be readily assessed using standard assays. For a review
 25 of the families of cytokines, see Mosmann and Coffman, *Ann. Rev. Immunol.* 7:145-173, 1989.

Preferred adjuvants for use in eliciting a predominantly Th1-type response include, for example, a combination of monophosphoryl lipid A, preferably 3-de-O-acylated monophosphoryl lipid A (3D-MPL), together with an aluminum salt. MPL

adjuvants are available from Ribl ImmunoChem Research Inc. (Hamilton, MT; *see* US Patent Nos. 4,436,727; 4,877,611; 4,866,034 and 4,912,094). Also preferred is AS-2 (SmithKline Beecham). CpG-containing oligonucleotides (in which the CpG dinucleotide is unmethylated) also induce a predominantly Th1 response. Such oligonucleotides are well known and are described, for example, in WO 96/02555. Another preferred adjuvant is a saponin, preferably QS21, which may be used alone or in combination with other adjuvants. For example, an enhanced system involves the combination of a monophosphoryl lipid A and saponin derivative, such as the combination of QS21 and 3D-MPL as described in WO 94/00153, or a less reactogenic composition where the QS21 is quenched with cholesterol, as described in WO 96/33739. Other preferred formulations comprises an oil-in-water emulsion and tocopherol. A particularly potent adjuvant formulation involving QS21, 3D-MPL and tocopherol in an oil-in-water emulsion is described in WO 95/17210. Any vaccine provided herein may be prepared using well known methods that result in a combination of antigen, immune response enhancer and a suitable carrier or excipient.

The compositions described herein may be administered as part of a sustained release formulation (*i.e.*, a formulation such as a capsule or sponge that effects a slow release of compound following administration). Such formulations may generally be prepared using well known technology and administered by, for example, oral, rectal or subcutaneous implantation, or by implantation at the desired target site. Sustained-release formulations may contain a polypeptide, polynucleotide or antibody dispersed in a carrier matrix and/or contained within a reservoir surrounded by a rate controlling membrane. Carriers for use within such formulations are biocompatible, and may also be biodegradable; preferably the formulation provides a relatively constant level of active component release. The amount of active compound contained within a sustained release formulation depends upon the site of implantation, the rate and expected duration of release and the nature of the condition to be treated or prevented.

Any of a variety of delivery vehicles may be employed within pharmaceutical compositions and vaccines to facilitate production of an antigen-specific

immune response that targets tumor cells. Delivery vehicles include antigen presenting cells (APCs), such as dendritic cells, macrophages, B cells, monocytes and other cells that may be engineered to be efficient APCs. Such cells may, but need not, be genetically modified to increase the capacity for presenting the antigen, to improve activation and/or maintenance of the T cell response, to have anti-tumor effects *per se* and/or to be immunologically compatible with the receiver (*i.e.*, matched HLA haplotype). APCs may generally be isolated from any of a variety of biological fluids and organs, including tumor and peritumoral tissues, and may be autologous, allogeneic, syngeneic or xenogeneic cells.

Certain preferred embodiments of the present invention use dendritic cells or progenitors thereof as antigen-presenting cells. Dendritic cells are highly potent APCs (Banchereau and Steinman, *Nature* 392:245-251, 1998) and have been shown to be effective as a physiological adjuvant for eliciting prophylactic or therapeutic antitumor immunity (*see* Timmerman and Levy, *Ann. Rev. Med.* 50:507-529, 1999). In general, dendritic cells may be identified based on their typical shape (stellate *in situ*, with marked cytoplasmic processes (dendrites) visible *in vitro*) and based on the lack of differentiation markers of B cells (CD19 and CD20), T cells (CD3), monocytes (CD14) and natural killer cells (CD56), as determined using standard assays. Dendritic cells may, of course, be engineered to express specific cell-surface receptors or ligands that are not commonly found on dendritic cells *in vivo* or *ex vivo*, and such modified dendritic cells are contemplated by the present invention. As an alternative to dendritic cells, secreted vesicles antigen-loaded dendritic cells (called exosomes) may be used within a vaccine (*see* Zitvogel et al., *Nature Med.* 4:594-600, 1998).

Dendritic cells and progenitors may be obtained from peripheral blood, bone marrow, tumor-infiltrating cells, peritumoral tissues-infiltrating cells, lymph nodes, spleen, skin, umbilical cord blood or any other suitable tissue or fluid. For example, dendritic cells may be differentiated *ex vivo* by adding a combination of cytokines such as GM-CSF, IL-4, IL-13 and/or TNF α to cultures of monocytes harvested from peripheral blood. Alternatively, CD34 positive cells harvested from peripheral blood, umbilical cord blood or bone marrow may be differentiated into dendritic cells by adding to the culture medium

combinations of GM-CSF, IL-3, TNF α , CD40 ligand, LPS, flt3 ligand and/or other compound(s) that induce maturation and proliferation of dendritic cells.

Dendritic cells are conveniently categorized as "immature" and "mature" cells, which allows a simple way to discriminate between two well characterized phenotypes. However, this nomenclature should not be construed to exclude all possible intermediate stages of differentiation. Immature dendritic cells are characterized as APC with a high capacity for antigen uptake and processing, which correlates with the high expression of Fc γ receptor, mannose receptor and DEC-205 marker. The mature phenotype is typically characterized by a lower expression of these markers, but a high expression of cell surface molecules responsible for T cell activation such as class I and class II MHC, adhesion molecules (*e.g.*, CD54 and CD11) and costimulatory molecules (*e.g.*, CD40, CD80 and CD86).

APCs may generally be transfected with a polynucleotide encoding a ovarian carcinoma antigen (or portion or other variant thereof) such that the antigen, or an immunogenic portion thereof, is expressed on the cell surface. Such transfection may take place *ex vivo*, and a composition or vaccine comprising such transfected cells may then be used for therapeutic purposes, as described herein. Alternatively, a gene delivery vehicle that targets a dendritic or other antigen presenting cell may be administered to a patient, resulting in transfection that occurs *in vivo*. *In vivo* and *ex vivo* transfection of dendritic cells, for example, may generally be performed using any methods known in the art, such as those described in WO 97/24447, or the gene gun approach described by Mahvi et al., *Immunology and cell Biology* 75:456-460, 1997. Antigen loading of dendritic cells may be achieved by incubating dendritic cells or progenitor cells with the polypeptide, DNA (naked or within a plasmid vector) or RNA; or with antigen-expressing recombinant bacterium or viruses (*e.g.*, vaccinia, fowlpox, adenovirus or lentivirus vectors). Prior to loading, the polypeptide may be covalently conjugated to an immunological partner that provides T cell help (*e.g.*, a carrier molecule). Alternatively, a dendritic cell may be pulsed with a non-conjugated immunological partner, separately or in the presence of the polypeptide.

CANCER THERAPY

In further aspects of the present invention, the compositions described herein may be used for immunotherapy of cancer, such as ovarian cancer. Within such methods, pharmaceutical compositions and vaccines are typically administered to a patient.

- 5 As used herein, a “patient” refers to any warm-blooded animal, preferably a human. A patient may or may not be afflicted with cancer. Accordingly, the above pharmaceutical compositions and vaccines may be used to prevent the development of a cancer or to treat a patient afflicted with a cancer. Within certain preferred embodiments, a patient is afflicted with ovarian cancer. Such cancer may be diagnosed using criteria generally accepted in the
- 10 art, including the presence of a malignant tumor. Pharmaceutical compositions and vaccines may be administered either prior to or following surgical removal of primary tumors and/or treatment such as administration of radiotherapy or conventional chemotherapeutic drugs.

- Within certain embodiments, immunotherapy may be active
- 15 immunotherapy, in which treatment relies on the *in vivo* stimulation of the endogenous host immune system to react against tumors with the administration of immuno response-modifying agents (such as tumor vaccines, bacterial adjuvants and/or cytokines).

- Within other embodiments, immunotherapy may be passive immunotherapy, in which treatment involves the delivery of agents with established tumor-immune
- 20 reactivity (such as effector cells or antibodies) that can directly or indirectly mediate antitumor effects and does not necessarily depend on an intact host immune system. Examples of effector cells include T lymphocytes (such as CD8⁺ cytotoxic T lymphocytes and CD4⁺ T-helper tumor-infiltrating lymphocytes), killer cells (such as Natural Killer cells and lymphokine-activated killer cells), B cells and antigen-presenting cells (such as
- 25 dendritic cells and macrophages) expressing a polypeptide provided herein. T cell receptors and antibody receptors specific for the polypeptides recited herein may be cloned, expressed and transferred into other vectors or effector cells for adoptive immunotherapy. The polypeptides provided herein may also be used to generate antibodies or anti-idiotypic

antibodies (as described above and in U.S. Patent No. 4,918,164) for passive immunotherapy.

Effector cells may generally be obtained in sufficient quantities for adoptive immunotherapy by growth *in vitro*, as described herein. Culture conditions for expanding single antigen-specific effector cells to several billion in number with retention of antigen recognition *in vivo* are well known in the art. Such *in vitro* culture conditions typically use intermittent stimulation with antigen, often in the presence of cytokines (such as IL-2) and non-dividing feeder cells. As noted above, immunoreactive polypeptides as provided herein may be used to rapidly expand antigen-specific T cell cultures in order to generate a sufficient number of cells for immunotherapy. In particular, antigen-presenting cells, such as dendritic, macrophage or B cells, may be pulsed with immunoreactive polypeptides or transfected with one or more polynucleotides using standard techniques well known in the art. For example, antigen-presenting cells can be transfected with a polynucleotide having a promoter appropriate for increasing expression in a recombinant virus or other expression system. Cultured effector cells for use in therapy must be able to grow and distribute widely, and to survive long term *in vivo*. Studies have shown that cultured effector cells can be induced to grow *in vivo* and to survive long term in substantial numbers by repeated stimulation with antigen supplemented with IL-2 (*see, for example, Cheever et al., Immunological Reviews 157:177, 1997*).

Alternatively, a vector expressing a polypeptide recited herein may be introduced into stem cells taken from a patient and clonally propagated *in vitro* for autologous transplant back into the same patient.

Routes and frequency of administration, as well as dosage, will vary from individual to individual, and may be readily established using standard techniques. In general, the pharmaceutical compositions and vaccines may be administered by injection (*e.g., intracutaneous, intramuscular, intravenous or subcutaneous*), intranasally (*e.g., by aspiration*), orally or in the bed of a resected tumor. Preferably, between 1 and 10 doses may be administered over a 52 week period. Preferably, 6 doses are administered, at intervals of 1 month, and booster vaccinations may be given periodically thereafter.

Alternate protocols may be appropriate for individual patients. A suitable dose is an amount of a compound that, when administered as described above, is capable of promoting an anti-tumor immune response, and is at least 10-50% above the basal (*i.e.*, untreated) level.. Such response can be monitored by measuring the anti-tumor antibodies

5 in a patient or by vaccine-dependent generation of cytolytic effector cells capable of killing the patient's tumor cells *in vitro*. Such vaccines should also be capable of causing an immune response that leads to an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial or longer disease-free survival) in vaccinated patients as compared to non-vaccinated patients. In general, for pharmaceutical compositions and

10 vaccines comprising one or more polypeptides, the amount of each polypeptide present in a dose ranges from about 100 µg to 5 mg per kg of host. Suitable dose sizes will vary with the size of the patient, but will typically range from about 0.1 mL to about 5 mL.

In general, an appropriate dosage and treatment regimen provides the active compound(s) in an amount sufficient to provide therapeutic and/or prophylactic benefit.

15 Such a response can be monitored by establishing an improved clinical outcome (*e.g.*, more frequent remissions, complete or partial, or longer disease-free survival) in treated patients as compared to non-treated patients. Increases in preexisting immune responses to an ovarian carcinoma antigen generally correlate with an improved clinical outcome. Such immune responses may generally be evaluated using standard proliferation, cytotoxicity or

20 cytokine assays, which may be performed using samples obtained from a patient before and after treatment.

SCREENS FOR IDENTIFYING SECRETED OVARIAN CARCINOMA ANTIGENS

The present invention provides methods for identifying secreted tumor

25 antigens. Within such methods, tumors are implanted into immunodeficient animals such as SCID mice and maintained for a time sufficient to permit secretion of tumor antigens into serum. In general, tumors may be implanted subcutaneously or within the gonadal fat pad of an immunodeficient animal and maintained for 1-9 months, preferably 1-4 months. Implantation may generally be performed as described in WO 97/18300. The serum

containing secreted antigens is then used to prepare antisera in immunocompetent mice, using standard techniques and as described herein. Briefly, 50-100 μ L of sera (pooled from three sets of immunodeficient mice, each set bearing a different SCID-derived human ovarian tumor) may be mixed 1:1 (vol:vol) with an appropriate adjuvant, such as RIBI-MPL or MPL + TDM (Sigma Chemical Co., St. Louis, MO) and injected intraperitoneally into syngeneic immunocompetent animals at monthly intervals for a total of 5 months. Antisera from animals immunized in such a manner may be obtained by drawing blood after the third, fourth and fifth immunizations. The resulting antiserum is generally pre-cleared of *E. coli* and phage antigens and used (generally following dilution, such as 1:200) in a serological expression screen.

The library is typically an expression library containing cDNAs from one or more tumors of the type that was implanted into SCID mice. This expression library may be prepared in any suitable vector, such as λ -screen (Novagen). cDNAs that encode a polypeptide that reacts with the antiserum may be identified using standard techniques, and sequenced. Such cDNA molecules may be further characterized to evaluate expression in tumor and normal tissue, and to evaluate antigen secretion in patients.

The methods provided herein have advantages over other methods for tumor antigen discovery. In particular, all antigens identified by such methods should be secreted or released through necrosis of the tumor cells. Such antigens may be present on the surface of tumor cells for an amount of time sufficient to permit targeting and killing by the immune system, following vaccination.

METHODS FOR DETECTING CANCER

In general, a cancer may be detected in a patient based on the presence of one or more ovarian carcinoma proteins and/or polynucleotides encoding such proteins in a biological sample (such as blood, sera, urine and/or tumor biopsies) obtained from the patient. In other words, such proteins may be used as markers to indicate the presence or absence of a cancer such as ovarian cancer. In addition, such proteins may be useful for the detection of other cancers. The binding agents provided herein generally permit detection

of the level of protein that binds to the agent in the biological sample. Polynucleotide primers and probes may be used to detect the level of mRNA encoding a tumor protein, which is also indicative of the presence or absence of a cancer. In general, an ovarian carcinoma-associated sequence should be present at a level that is at least three fold higher
 5 in tumor tissue than in normal tissue

There are a variety of assay formats known to those of ordinary skill in the art for using a binding agent to detect polypeptide markers in a sample. *See, e.g.,* Harlow and Lane, *Antibodies: A Laboratory Manual*, Cold Spring Harbor Laboratory, 1988. In general, the presence or absence of a cancer in a patient may be determined by (a)
 10 contacting a biological sample obtained from a patient with a binding agent; (b) detecting in the sample a level of polypeptide that binds to the binding agent; and (c) comparing the level of polypeptide with a predetermined cut-off value.

In a preferred embodiment, the assay involves the use of binding agent immobilized on a solid support to bind to and remove the polypeptide from the remainder
 15 of the sample. The bound polypeptide may then be detected using a detection reagent that contains a reporter group and specifically binds to the binding agent/polypeptide complex. Such detection reagents may comprise, for example, a binding agent that specifically binds to the polypeptide or an antibody or other agent that specifically binds to the binding agent, such as an anti-immunoglobulin, protein G, protein A or a lectin. Alternatively, a
 20 competitive assay may be utilized, in which a polypeptide is labeled with a reporter group and allowed to bind to the immobilized binding agent after incubation of the binding agent with the sample. The extent to which components of the sample inhibit the binding of the labeled polypeptide to the binding agent is indicative of the reactivity of the sample with the immobilized binding agent. Suitable polypeptides for use within such assays include
 25 full length ovarian carcinoma proteins and portions thereof to which the binding agent binds, as described above.

The solid support may be any material known to those of ordinary skill in the art to which the tumor protein may be attached. For example, the solid support may be a test well in a microtiter plate or a nitrocellulose or other suitable membrane.

Alternatively, the support may be a bead or disc, such as glass, fiberglass, latex or a plastic material such as polystyrene or polyvinylchloride. The support may also be a magnetic particle or a fiber optic sensor, such as those disclosed, for example, in U.S. Patent No. 5,359,681. The binding agent may be immobilized on the solid support using a variety of techniques known to those of skill in the art, which are amply described in the patent and scientific literature. In the context of the present invention, the term "immobilization" refers to both noncovalent association, such as adsorption, and covalent attachment (which may be a direct linkage between the agent and functional groups on the support or may be a linkage by way of a cross-linking agent). Immobilization by adsorption to a well in a microtiter plate or to a membrane is preferred. In such cases, adsorption may be achieved by contacting the binding agent, in a suitable buffer, with the solid support for a suitable amount of time. The contact time varies with temperature, but is typically between about 1 hour and about 1 day. In general, contacting a well of a plastic microtiter plate (such as polystyrene or polyvinylchloride) with an amount of binding agent ranging from about 10 ng to about 10 μ g, and preferably about 100 ng to about 1 μ g, is sufficient to immobilize an adequate amount of binding agent.

Covalent attachment of binding agent to a solid support may generally be achieved by first reacting the support with a bifunctional reagent that will react with both the support and a functional group, such as a hydroxyl or amino group, on the binding agent. For example, the binding agent may be covalently attached to supports having an appropriate polymer coating using benzoquinone or by condensation of an aldehyde group on the support with an amine and an active hydrogen on the binding partner (*see, e.g.*, Pierce Immunotechnology Catalog and Handbook, 1991, at A12-A13).

In certain embodiments, the assay is a two-antibody sandwich assay. This assay may be performed by first contacting an antibody that has been immobilized on a solid support, commonly the well of a microtiter plate, with the sample, such that polypeptides within the sample are allowed to bind to the immobilized antibody. Unbound sample is then removed from the immobilized polypeptide-antibody complexes and a detection reagent (preferably a second antibody capable of binding to a different site on the

polypeptide) containing a reporter group is added. The amount of detection reagent that remains bound to the solid support is then determined using a method appropriate for the specific reporter group.

More specifically, once the antibody is immobilized on the support as described above, the remaining protein binding sites on the support are typically blocked. Any suitable blocking agent known to those of ordinary skill in the art, such as bovine serum albumin or Tween 20™ (Sigma Chemical Co., St. Louis, MO). The immobilized antibody is then incubated with the sample, and polypeptide is allowed to bind to the antibody. The sample may be diluted with a suitable diluent, such as phosphate-buffered saline (PBS) prior to incubation. In general, an appropriate contact time (*i.e.*, incubation time) is a period of time that is sufficient to detect the presence of polypeptide within a sample obtained from an individual with ovarian cancer. Preferably, the contact time is sufficient to achieve a level of binding that is at least about 95% of that achieved at equilibrium between bound and unbound polypeptide. Those of ordinary skill in the art will recognize that the time necessary to achieve equilibrium may be readily determined by assaying the level of binding that occurs over a period of time. At room temperature, an incubation time of about 30 minutes is generally sufficient.

Unbound sample may then be removed by washing the solid support with an appropriate buffer, such as PBS containing 0.1% Tween 20™. The second antibody, which contains a reporter group, may then be added to the solid support. Preferred reporter groups include those groups recited above.

The detection reagent is then incubated with the immobilized antibody-polypeptide complex for an amount of time sufficient to detect the bound polypeptide. An appropriate amount of time may generally be determined by assaying the level of binding that occurs over a period of time. Unbound detection reagent is then removed and bound detection reagent is detected using the reporter group. The method employed for detecting the reporter group depends upon the nature of the reporter group. For radioactive groups, scintillation counting or autoradiographic methods are generally appropriate. Spectroscopic methods may be used to detect dyes, luminescent groups and fluorescent groups. Biotin

may be detected using avidin, coupled to a different reporter group (commonly a radioactive or fluorescent group or an enzyme). Enzyme reporter groups may generally be detected by the addition of substrate (generally for a specific period of time), followed by spectroscopic or other analysis of the reaction products.

5 To determine the presence or absence of a cancer, such as ovarian cancer, the signal detected from the reporter group that remains bound to the solid support is generally compared to a signal that corresponds to a predetermined cut-off value. In one preferred embodiment, the cut-off value for the detection of a cancer is the average mean signal obtained when the immobilized antibody is incubated with samples from patients
10 without the cancer. In general, a sample generating a signal that is three standard deviations above the predetermined cut-off value is considered positive for the cancer. In an alternate preferred embodiment, the cut-off value is determined using a Receiver Operator Curve, according to the method of Sackett et al., *Clinical Epidemiology: A Basic Science for Clinical Medicine*, Little Brown and Co., 1985, p. 106-7. Briefly, in this
15 embodiment, the cut-off value may be determined from a plot of pairs of true positive rates (i.e., sensitivity) and false positive rates (100%-specificity) that correspond to each possible cut-off value for the diagnostic test result. The cut-off value on the plot that is the closest to the upper left-hand corner (i.e., the value that encloses the largest area) is the most accurate cut-off value, and a sample generating a signal that is higher than the cut-off value
20 determined by this method may be considered positive. Alternatively, the cut-off value may be shifted to the left along the plot, to minimize the false positive rate, or to the right, to minimize the false negative rate. In general, a sample generating a signal that is higher than the cut-off value determined by this method is considered positive for a cancer.

In a related embodiment, the assay is performed in a flow-through or strip
25 test format, wherein the binding agent is immobilized on a membrane, such as nitrocellulose. In the flow-through test, polypeptides within the sample bind to the immobilized binding agent as the sample passes through the membrane. A second, labeled binding agent then binds to the binding agent-polypeptide complex as a solution containing the second binding agent flows through the membrane. The detection of bound second

binding agent may then be performed as described above. In the strip test format, one end of the membrane to which binding agent is bound is immersed in a solution containing the sample. The sample migrates along the membrane through a region containing second binding agent and to the area of immobilized binding agent. Concentration of second binding agent at the area of immobilized antibody indicates the presence of a cancer. Typically, the concentration of second binding agent at that site generates a pattern, such as a line, that can be read visually. The absence of such a pattern indicates a negative result. In general, the amount of binding agent immobilized on the membrane is selected to generate a visually discernible pattern when the biological sample contains a level of polypeptide that would be sufficient to generate a positive signal in the two-antibody sandwich assay, in the format discussed above. Preferred binding agents for use in such assays are antibodies and antigen-binding fragments thereof. Preferably, the amount of antibody immobilized on the membrane ranges from about 25 ng to about 1 μ g, and more preferably from about 50 ng to about 500 ng. Such tests can typically be performed with a very small amount of biological sample.

Of course, numerous other assay protocols exist that are suitable for use with the tumor proteins or binding agents of the present invention. The above descriptions are intended to be exemplary only. For example, it will be apparent to those of ordinary skill in the art that the above protocols may be readily modified to use ovarian carcinoma polypeptides to detect antibodies that bind to such polypeptides in a biological sample. The detection of such ovarian carcinoma protein specific antibodies may correlate with the presence of a cancer.

A cancer may also, or alternatively, be detected based on the presence of T cells that specifically react with an ovarian carcinoma protein in a biological sample. Within certain methods, a biological sample comprising CD4⁺ and/or CD8⁺ T cells isolated from a patient is incubated with an ovarian carcinoma protein, a polynucleotide encoding such a polypeptide and/or an APC that expresses at least an immunogenic portion of such a polypeptide, and the presence or absence of specific activation of the T cells is detected. Suitable biological samples include, but are not limited to, isolated T cells. For example, T

cells may be isolated from a patient by routine techniques (such as by Ficoll/Hypaque density gradient centrifugation of peripheral blood lymphocytes). T cells may be incubated *in vitro* for 2-9 days (typically 4 days) at 37°C with an ovarian carcinoma protein (e.g., 5 - 25 µg/ml). It may be desirable to incubate another aliquot of a T cell sample in the absence
 5 of ovarian carcinoma protein to serve as a control. For CD4⁺ T cells, activation is preferably detected by evaluating proliferation of the T cells. For CD8⁺ T cells, activation is preferably detected by evaluating cytolytic activity. A level of proliferation that is at least two fold greater and/or a level of cytolytic activity that is at least 20% greater than in disease-free patients indicates the presence of a cancer in the patient.

10 As noted above, a cancer may also, or alternatively, be detected based on the level of mRNA encoding an ovarian carcinoma protein in a biological sample. For example, at least two oligonucleotide primers may be employed in a polymerase chain reaction (PCR) based assay to amplify a portion of an ovarian carcinoma protein cDNA derived from a biological sample, wherein at least one of the oligonucleotide primers is
 15 specific for (*i.e.*, hybridizes to) a polynucleotide encoding the ovarian carcinoma protein. The amplified cDNA is then separated and detected using techniques well known in the art, such as gel electrophoresis. Similarly, oligonucleotide probes that specifically hybridize to a polynucleotide encoding an ovarian carcinoma protein may be used in a hybridization assay to detect the presence of polynucleotide encoding the tumor protein in a biological
 20 sample.

To permit hybridization under assay conditions, oligonucleotide primers and probes should comprise an oligonucleotide sequence that has at least about 60%, preferably at least about 75% and more preferably at least about 90%, identity to a portion of a polynucleotide encoding an ovarian carcinoma protein that is at least 10 nucleotides, and
 25 preferably at least 20 nucleotides, in length. Preferably, oligonucleotide primers and/or probes hybridize to a polynucleotide encoding a polypeptide described herein under moderately stringent conditions, as defined above. Oligonucleotide primers and/or probes which may be usefully employed in the diagnostic methods described herein preferably are at least 10-40 nucleotides in length. In a preferred embodiment, the oligonucleotide

primers comprise at least 10 contiguous nucleotides, more preferably at least 15 contiguous nucleotides, of a DNA molecule having a sequence provided herein. Techniques for both PCR based assays and hybridization assays are well known in the art (*see*, for example, Mullis et al., *Cold Spring Harbor Symp. Quant. Biol.*, 51:263, 1987; Erlich ed., *PCR Technology*, Stockton Press, NY, 1989).

One preferred assay employs RT-PCR, in which PCR is applied in conjunction with reverse transcription. Typically, RNA is extracted from a biological sample such as a biopsy tissue and is reverse transcribed to produce cDNA molecules. PCR amplification using at least one specific primer generates a cDNA molecule, which may be separated and visualized using, for example, gel electrophoresis. Amplification may be performed on biological samples taken from a test patient and from an individual who is not afflicted with a cancer. The amplification reaction may be performed on several dilutions of cDNA spanning two orders of magnitude. A two-fold or greater increase in expression in several dilutions of the test patient sample as compared to the same dilutions of the non-cancerous sample is typically considered positive.

In another embodiment, ovarian carcinoma proteins and polynucleotides encoding such proteins may be used as markers for monitoring the progression of cancer. In this embodiment, assays as described above for the diagnosis of a cancer may be performed over time, and the change in the level of reactive polypeptide(s) evaluated. For example, the assays may be performed every 24-72 hours for a period of 6 months to 1 year, and thereafter performed as needed. In general, a cancer is progressing in those patients in whom the level of polypeptide detected by the binding agent increases over time. In contrast, the cancer is not progressing when the level of reactive polypeptide either remains constant or decreases with time.

Certain *in vivo* diagnostic assays may be performed directly on a tumor. One such assay involves contacting tumor cells with a binding agent. The bound binding agent may then be detected directly or indirectly via a reporter group. Such binding agents may also be used in histological applications. Alternatively, polynucleotide probes may be used within such applications.

As noted above, to improve sensitivity, multiple ovarian carcinoma protein markers may be assayed within a given sample. It will be apparent that binding agents specific for different proteins provided herein may be combined within a single assay. Further, multiple primers or probes may be used concurrently. The selection of tumor protein markers may be based on routine experiments to determine combinations that results in optimal sensitivity. In addition, or alternatively, assays for tumor proteins provided herein may be combined with assays for other known tumor antigens.

DIAGNOSTIC KITS

The present invention further provides kits for use within any of the above diagnostic methods. Such kits typically comprise two or more components necessary for performing a diagnostic assay. Components may be compounds, reagents, containers and/or equipment. For example, one container within a kit may contain a monoclonal antibody or fragment thereof that specifically binds to an ovarian carcinoma protein. Such antibodies or fragments may be provided attached to a support material, as described above. One or more additional containers may enclose elements, such as reagents or buffers, to be used in the assay. Such kits may also, or alternatively, contain a detection reagent as described above that contains a reporter group suitable for direct or indirect detection of antibody binding.

Alternatively, a kit may be designed to detect the level of mRNA encoding an ovarian carcinoma protein in a biological sample. Such kits generally comprise at least one oligonucleotide probe or primer, as described above, that hybridizes to a polynucleotide encoding an ovarian carcinoma protein. Such an oligonucleotide may be used, for example, within a PCR or hybridization assay. Additional components that may be present within such kits include a second oligonucleotide and/or a diagnostic reagent or container to facilitate the detection of a polynucleotide encoding an ovarian carcinoma protein.

The following Examples are offered by way of illustration and not by way of limitation.

EXAMPLES

Example 1

Identification of Representative Ovarian Carcinoma Protein cDNAs

5

This Example illustrates the identification of cDNA molecules encoding ovarian carcinoma proteins.

Anti-SCID mouse sera (generated against sera from SCID mice carrying late passage ovarian carcinoma) was pre-cleared of *E. coli* and phage antigens and used at a
 10 1:200 dilution in a serological expression screen. The library screened was made from a SCID-derived human ovarian tumor (OV9334) using a directional RH oligo(dT) priming cDNA library construction kit and the λ Screen vector (Novagen). A bacteriophage lambda screen was employed. Approximately 400,000 pfu of the amplified OV9334 library were screened.

15 196 positive clones were isolated. Certain sequences that appear to be novel are provided in Figures 1A-1S and SEQ ID NOs:1 to 71. Three complete insert sequences are shown in Figures 2A-2C (SEQ ID NOs:72 to 74). Other clones having known sequences are presented in Figures 15A-15EEE (SEQ ID NOs:82 to 310). Database searches identified the following sequences that were substantially identical to the
 20 sequences presented in Figures 15A-15EEE.

These clones were further characterized using microarray technology to determine mRNA expression levels in a variety of tumor and normal tissues. Such analyses were performed using a Synteni (Palo Alto, CA) microarray, according to the manufacturer's instructions. PCR amplification products were arrayed on slides, with each
 25 product occupying a unique location in the array. mRNA was extracted from the tissue sample to be tested, reverse transcribed and fluorescent-labeled cDNA probes were generated. The microarrays were probed with the labeled cDNA probes and the slides were scanned to measure fluorescence intensity. Data was analyzed using Synteni's provided

GEMtools software. The results for one clone (13695, also referred to as O8E) are shown in Figure 3.

Example 2

5 Identification of Ovarian Carcinoma cDNAs using Microarray Technology

This Example illustrates the identification of ovarian carcinoma polynucleotides by PCR subtraction and microarray analysis. Microarrays of cDNAs were analyzed for ovarian tumor-specific expression using a Synteni (Palo Alto, CA) microarray, according to the manufacturer's instructions (and essentially as described by Schena et al.,
10 *Proc. Natl. Acad. Sci. USA* 93:10614-10619, 1996 and Heller et al., *Proc. Natl. Acad. Sci. USA* 94:2150-2155, 1997).

A PCR subtraction was performed using a tester comprising cDNA of four ovarian tumors (three of which were metastatic tumors) and a driver of cDNA from five
15 normal tissues (adrenal gland, lung, pancreas, spleen and brain). cDNA fragments recovered from this subtraction were subjected to DNA microarray analysis where the fragments were PCR amplified, adhered to chips and hybridized with fluorescently labeled probes derived from mRNAs of human ovarian tumors and a variety of normal human tissues. In this analysis, the slides were scanned and the fluorescence intensity was
20 measured, and the data were analyzed using Synteni's GEMtools software. In general, sequences showing at least a 5-fold increase in expression in tumor cells (relative to normal cells) were considered ovarian tumor antigens. The fluorescent results were analyzed and clones that displayed increased expression in ovarian tumors were further characterized by DNA sequencing and database searches to determine the novelty of the sequences.

25 Using such assays, an ovarian tumor antigen was identified that is a splice fusion between the human T-cell leukemia virus type I oncoprotein TAX (*see* Jin et al., *Cell* 93:81-91, 1998) and an extracellular matrix protein called osteonectin. A splice junction sequence exists at the fusion point. The sequence of this clone is presented in

Figure 4 and SEQ ID NO:75. Osteonectin, unspliced and unaltered, was also identified from such assays independently.

Further clones identified by this method are referred to herein as 3f, 6b, 8e, 8h, 12c and 12h. Sequences of these clones are shown in Figures 5 to 9 and SEQ ID NOs:76 to 81. Microarray analyses were performed as described above, and are presented in Figures 10 to 14. A full length sequence encompassing clones 3f, 6b, 8e and 12h was obtained by screening an ovarian tumor (SCID-derived) cDNA library. This 2996 base pair sequence (designated O772P) is presented in SEQ ID NO:311, and the encoded 914 amino acid protein sequence is shown in SEQ ID NO:312. PSORT analysis indicates a Type 1a transmembrane protein localized to the plasma membrane.

In addition to certain of the sequences described above, this screen identified the following sequences:

Sequence	Comments
OV4vG11 (SEQ ID NO:313)	human clone 1119D9 on chromosome 20p12
OV4vB11 (SEQ ID NO:314)	human UWGC:y14c094 from chromosome 6p21
OV4vD9 (SEQ ID NO:315)	human clone 1049G16 chromosome 20q12-13.2
OV4vD5 (SEQ ID NO:316)	human KIAA0014 gene
OV4vC2 (SEQ ID NO:317)	human KIAA0084 gene
OV4vF3 (SEQ ID NO:318)	human chromosome 19 cosmid R31167
OV4VC1 (SEQ ID NO:319)	novel
OV4vH3 (SEQ ID NO:320)	novel
OV4vD2 (SEQ ID NO:321)	novel
O815P (SEQ ID NO:322)	novel
OV4vC12 (SEQ ID NO:323)	novel
OV4vA4 (SEQ ID NO:324)	novel
OV4vA3 (SEQ ID NO:325)	novel
OV4v2A5 (SEQ ID NO:326)	novel
O819P (SEQ ID NO:327)	novel
O818P (SEQ ID NO:328)	novel
O817P (SEQ ID NO:329)	novel

Sequence	Comments
O816P (SEQ ID NO:330)	novel
Ov4vC5 (SEQ ID NO:331)	novel
21721 (SEQ ID NO:332)	human lumican
21719 (SEQ ID NO:333)	human retinoic acid-binding protein II
21717 (SEQ ID NO:334)	human26S proteasome ATPase subunit
21654 (SEQ ID NO:335)	human copine I
21627 (SEQ ID NO:336)	human neuron specific gamma-2 enolase
21623 (SEQ ID NO:337)	human geranylgeranyl transferase II
21621 (SEQ ID NO:338)	human cyclin-dependent protein kinase
21616 (SEQ ID NO:339)	human prepro-megakaryocyte potentiating factor
21612 (SEQ ID NO:340)	human UPH1
21558 (SEQ ID NO:341)	human RalGDS-like 2 (RGL2)
21555 (SEQ ID NO:342)	human autoantigen P542
21548 (SEQ ID NO:343)	human actin-related protein (ARP2)
21462 (SEQ ID NO:344)	human huntingtin interacting protein
21441 (SEQ ID NO:345)	human 90K product (tumor associated antigen)
21439 (SEQ ID NO:346)	human guanine nucleotide regulator protein (tim1)
21438 (SEQ ID NO:347)	human Ku autoimmune (p70/p80) antigen
21237 (SEQ ID NO:348)	human S-laminin
21436 (SEQ ID NO:349)	human ribophorin I
21435 (SEQ ID NO:350)	human cytoplasmic chaperonin hTRiC5
21425 (SEQ ID NO:351)	humanEMX2
21423 (SEQ ID NO:352)	human p87/p89 gene
21419 (SEQ ID NO:353)	human HPBR11-7
21252 (SEQ ID NO:354)	human T1-227H
21251 (SEQ ID NO:355)	human cullin I
21247 (SEQ ID NO:356)	kunitz type protease inhibitor (KOP)
21244-1 (SEQ ID NO:357)	human protein tyrosine phosphatase receptor F (PTPRF)
21718 (SEQ ID NO:358)	human LTR repeat
OV2-90 (SEQ ID NO:359)	novel
Human zinc finger (SEQ ID NO:360)	

Sequence	Comments
Human polyA binding protein (SEQ ID NO:361)	
Human pleiotrophin (SEQ ID NO:362)	
Human PAC clone 278C19 (SEQ ID NO:363)	
Human LLRep3 (SEQ ID NO:364)	
Human Kunitz type protease inhib (SEQ ID NO:365)	
Human KIAA0106 gene (SEQ ID NO:366)	
Human keratin (SEQ ID NO:367)	
Human HIV-1TAR (SEQ ID NO:368)	
Human glia derived nexin (SEQ ID NO:369)	
Human fibronectin (SEQ ID NO:370)	
Human ECMproBM40 (SEQ ID NO:371)	
Human collagen (SEQ ID NO:372)	
Human alpha enolase (SEQ ID NO:373)	
Human aldolase (SEQ ID NO:374)	
Human transf growth factor BIG H3 (SEQ ID NO:375)	
Human SPARC osteonectin (SEQ ID NO:376)	
Human SLP1 leucocyte protease (SEQ ID NO:377)	
Human mitochondrial ATP synth (SEQ ID NO:378)	
Human DNA seq clone 461P17 (SEQ ID NO:379)	
Human dbpB pro Y box (SEQ ID NO:380)	
Human 40 kDa keratin (SEQ ID NO:381)	
Human arginosuccinate synth (SEQ ID NO:382)	
Human acidic ribosomal phosphoprotein (SEQ ID NO:383)	
Human colon carcinoma laminin binding pro (SEQ ID NO:384)	

This screen further identified multiple forms of the clone O772P, referred to herein as 21013, 21003 and 21008. PSORT analysis indicates that 21003 (SEQ ID NO:386; translated as SEQ ID NO:389) and 21008 (SEQ ID NO:387; translated as SEQ ID NO:390) represent Type 1a transmembrane protein forms of O772P. 21013 (SEQ ID NO:385; translated as SEQ ID NO:388) appears to be a truncated form of the protein and is predicted by PSORT analysis to be a secreted protein.

Additional sequence analysis resulted in a full length clone for O8E (2627 bp, which agrees with the message size observed by Northern analysis; SEQ ID NO:391). This nucleotide sequence was obtained as follows: the original O8E sequence (OrigO8Econs) was found to overlap by 33 nucleotides with a sequence from an EST clone
 5 (IMAGE#1987589). This clone provided 1042 additional nucleotides upstream of the original O8E sequence. The link between the EST and O8E was confirmed by sequencing multiple PCR fragments generated from an ovary primary tumor library using primers to the unique EST and the O8E sequence (ESTxO8EPCR). Full length status was further indicated when anchored PCR from the ovary tumor library gave several clones
 10 (AnchoredPCR cons) that all terminated upstream of the putative start methionine, but failed to yield any additional sequence information. Figure 16 presents a diagram that illustrates the location of each partial sequence within the full length O8E sequence.

Two protein sequences may be translated from the full length O8E. For "a" (SEQ ID NO:393) begins with a putative start methionine. A second form "b" (SEQ ID
 15 NO:392) includes 27 additional upstream residues to the 5' end of the nucleotide sequence.

Example 3

This example discloses the identification and characterization of antibody
 20 epitopes recognized by the O8E polyclonal anti-sera.

Rabbit anti-sera was raised against E. coli derived O8E recombinant protein and tested for antibody epitope recognition against 20 or 21 mer peptides that correspond to the O8E amino acid sequence. Peptides spanning amino acid regions 31 to 65, 76 to 110, 136 to 200 and 226 to 245 of the full length O8E protein were recognized by an acid eluted
 25 peak and/or a salt eluted peak from affinity purified anti-O8E sera. Thus, the corresponding amino acid sequences of the above peptides constitute the antibody epitopes recognized by affinity purified anti-O8E antibodies.

ELISA analysis of anti-O8E rabbit sera is shown in Figure 23, and ELISA analysis of affinity purified rabbit anti-O8E polyclonal antibody is shown in Figure 24.

For epitope mapping, 20 or 21 mer peptides corresponding to the O8E protein were synthesized. For antibody affinity purification, rabbit anti-O8E sera was run over an O8E-sepharose column, then antibody was eluted with a salt buffer containing 0.5 M NaCl and 20 mM PO₄, followed by an acid elution step using 0.2 M Glycine, pH 2.3.

5 Purified antibody was neutralized by the addition of 1M Tris, pH 8 and buffer exchanged into phosphate buffered saline (PBS). For enzyme linked immunosorbant assay (ELISA) analysis, O8E peptides and O8E recombinant protein were coated onto 96 well flat bottom plates at 2 µg/ml for 2 hours at room temperature (RT). Plates were then washed 5 times with PBS + 0.1 % Tween 20 and blocked with PBS + 1 % bovine serum albumin (BSA) for

10 1 hour. Affinity purified anti-O8E antibody, either an acid or salt eluted fraction, was then added to the wells at 1 µg/ml and incubated at RT for 1 hr. Plates were again washed, followed by the addition of donkey anti-rabbit-Ig-horseradish peroxidase (HRP) antibody for 1 hour at RT. Plates were washed, then developed by the addition of the chromagenic substrate 3, 3', 5, 5'-tetramethylbenzidine (TMB) (described by Bos *et al.*, *J. of*

15 *Immunoassay* 2:187-204 (1981); available from Sigma (St. Louis, MO)). The reaction was incubated 15 minutes at RT and then stopped by the addition of 1 N H₂SO₄. Plates were read at an optical density of 450 (OD450) in an automated plate reader. The sequences of peptides corresponding to the OE8 antibody epitopes are disclosed herein as SEQ ID NOs: 394-415. Antibody epitopes recognized by the O8E polyclonal anti-sera are disclosed

20 herein in Figure 17.

Example 4

This example discloses IHC analysis of O8E expression in ovarian cancer

25 tissue samples.

For immunohistochemistry studies, paraffin-embedded formalin fixed ovarian cancer tissue was sliced into 8 micron sections. Steam heat induced epitope retrieval (SHIER) in 0.1 M sodium citrate buffer (pH 6.0) was used for optimal staining conditions. Sections were incubated with 10% serum/PBS for 5 minutes. Primary antibody

(anti-O8E rabbit affinity purified polyclonal antibody) was added to each section for 25 min followed by a 25 min incubation with an anti-rabbit biotinylated antibody. Endogenous peroxidase activity was blocked by three 1.5 min incubations with hydrogen peroxidase. The avidin biotin complex/horse radish peroxidase system was used along with DAB chromogen to visualize antigen expression. Slides were counterstained with hematoxylin. One (papillary serous carcinoma) of six ovarian cancer tissue sections displayed O8E immunoreactivity. O8E expression was localized to the plasma membrane.

Six ovarian cancer tissues were analyzed with the anti-O8E rabbit polyclonal antibody. One (papillary serous carcinoma) of six ovarian cancer tissue samples stained positive for O8E expression. O8E expression was localized to the surface membrane.

Example 5

This example discloses O8E peptides that are predicted to bind HLA-A2 and to be immunogenic for CD8 T cell responses in humans.

Potential HLA-A2 binding peptides of O8E were predicted by using the full-length open-reading frame (ORF) from O8E and running it through "Episeek," a program used to predict MHC binding peptides. The program used is based on the algorithm published by Parker, K.C. *et al.*, *J. Immunol.* 152(1):163-175 (1994) (incorporated by reference herein in its entirety). 10-mer and 9-mer peptides predicted to bind HLA-A2 are disclosed herein as SEQ ID NOs: 416-435 and SEQ ID NOs: 436-455, respectively.

Example 6

This example discloses O8E cell surface expression measured by fluorescence activated cell sorting.

For FACS analysis, cells were washed with ice cold staining buffer (PBS/1% BSA/azide). Next, the cells were incubated for 30 minutes on ice with 10

micrograms/ml of affinity purified rabbit anti-B305D polyclonal antibody. The cells were washed 3 times with staining buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig (H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing prodium iodide, a vital stain that allows for identification of permeable cells, and analyzed by FACS. O8E surface expression was confirmed on SKBR3 breast cancer cells and HEK293 cells that stably overexpress the cDNA for O8E. Neither MB415 cells nor HEK293 cells stably transfected with a control irrelevant plasmid DNA showed surface expression of O8E (Figures 18 and 19).

10

Example 7

This example further evaluates the expression and surface localization of O8E.

15

For expression and purification of antigen used for immunization, O8E expressed in an E. coli recombinant expression system was grown overnight in LB Broth with the appropriate antibiotics at 37°C in a shaking incubator. The next morning, 10 ml of the overnight culture was added to 500 ml of 2x YT plus appropriate antibiotics in a 2L-baffled Erlenmeyer flask. When the Optical Density (at 560 nanometers) of the culture reached 0.4-0.6 the cells were induced with IPTG (1 mM). 4 hours after induction with IPTG the cells were harvested by centrifugation. The cells were then washed with phosphate buffered saline and centrifuged again. The supernatant was discarded and the cells were either frozen for future use or immediately processed. Twenty milliliters of lysis buffer was added to the cell pellets and vortexed. To break open the E. coli cells, this mixture was then run through the French Press at a pressure of 16,000 psi. The cells were then centrifuged again and the supernatant and pellet were checked by SDS-PAGE for the partitioning of the recombinant protein. For protein that localized to the cell pellet, the pellet was resuspended in 10 mM Tris pH 8.0 , 1% CHAPS and the inclusion body pellet was washed and centrifuged again. This procedure was repeated twice more. The washed

25

inclusion body pellet was solubilized with either 8 M urea or 6 M guanidine HCl containing 10 mM Tris pH 8.0 plus 10 mM imidazole. The solubilized protein was added to 5 ml of nickel-chelate resin (Qiagen) and incubated for 45 min to 1 hour at room temperature with continuous agitation. After incubation, the resin and protein mixture were poured through a disposable column and the flow through was collected. The column was then washed with 10-20 column volumes of the solubilization buffer. The antigen was then eluted from the column using 8M urea, 10 mM tris pH 8.0 and 300 mM imidazole and collected in 3 ml fractions. A SDS-PAGE gel was run to determine which fractions to pool for further purification. As a final purification step, a strong anion exchange resin such as Hi-Prep Q (Biorad) was equilibrated with the appropriate buffer and the pooled fractions from above were loaded onto the column. Each antigen was eluted off of the column with an increasing salt gradient. Fractions were collected as the column was run and another SDS-PAGE gel was run to determine which fractions from the column to pool. The pooled fractions were dialyzed against 10 mM Tris pH 8.0. This material was then evaluated for acceptable purity as determined by SDS-PAGE or HPLC, concentration as determined by Lowry assay or Amino Acid Analysis, identity as determined by amino terminal protein sequence, and endotoxin level as determined by the Limulus (LAL) assay. The proteins were then vialled after filtration through a 0.22 micron filter and the antigens were frozen until needed for immunization.

For generation of polyclonal anti-sera, 400 micrograms of each prostate antigen was combined with 100 micrograms of muramyl dipeptide (MDP). Equal volume of Incomplete Freund's Adjuvant (IFA) was added and then mixed. Every four weeks animals were boosted with 100 micrograms of antigen mixed with an equal volume of IFA. Seven days following each boost the animal was bled. Sera was generated by incubating the blood at 4°C for 12-24 hours followed by centrifugation.

For characterization of polyclonal antisera, 96 well plates were coated with antigen by incubating with 50 microliters (typically 1 microgram) at 4 C for 20 hrs. 250 microliters of BSA blocking buffer was added to the wells and incubated at RT for 2 hrs. Plates were washed 6 times with PBS/0.01% tween. Anti-O8E rabbit sera or affinity

purified anti-O8e antibody was diluted in PBS. Fifty microliters of diluted antibody was added to each well and incubated at RT for 30 min. Plates were washed as described above before 50 microliters of goat anti-rabbit horse radish peroxidase (HRP) at a 1:10000 dilution was added and incubated at RT for 30 min. Plates were washed as described above and 100 microliters of TMB microwell Peroxidase Substrate was added to each well. Following a 15 minute incubation in the dark at room temperature the colorimetric reaction was stopped with 100 microliters of 1N H₂SO₄ and read immediately at 450 nm. All polyclonal antibodies showed immunoreactivity to the O8E antigen.

For recombinant expression in mammalian HEK293 cells, full length O8E cDNA was subcloned into the mammalian expression vectors pcDNA3.1+ and pCEP4 (Invitrogen) which were modified to contain His and FLAG epitope tags, respectively. These constructs were transfected into HEK293 cells (ATCC) using Fugene 6 reagent (Roche). Briefly, HEK293 cells were plated at a density of 100,000 cells/ml in DMEM (Gibco) containing 10% FBS (Hyclone) and grown overnight. The following day, 2 ul of Fugene6 was added to 100 ul of DMEM containing no FBS and incubated for 15 minutes at room temperature. The Fugene6/DMEM mixture was then added to 1ug of O8E/pCEP4 or O8E/pcDNA3.1 plasmid DNA and incubated for 15 minutes at room temperature. The Fugene/DNA mix was then added to the HEK293 cells and incubated for 48-72 hrs at 37°C with 7% CO₂. Cells were rinsed with PBS then collected and pelleted by centrifugation. For Western blot analysis, whole cell lysates were generated by incubating the cells in Triton-X100 containing lysis buffer for 30 minutes on ice. Lysates were then cleared by centrifugation at 10,000rpm for 5 minutes at 4 C. Samples were diluted with SDS-PAGE loading buffer containing beta-mercaptoethanol, then boiled for 10 minutes prior to loading the SDS-PAGE gel. Protein was transferred to nitrocellulose and probed using anti-O8E rabbit polyclonal sera #2333L at a dilution of 1:750. The blot was revealed with a goat anti-rabbit Ig coupled to HRP followed by incubation in ECL substrate.

For FACS analysis, cells were washed further with ice cold staining buffer (PBS+1%BSA+Azide). Next, the cells were incubated for 30 minutes on ice with 10ug/ml of Protein A purified anti-O8E polyclonal sera. The cells were washed 3 times with staining

buffer and then incubated with a 1:100 dilution of a goat anti-rabbit Ig(H+L)-FITC reagent (Southern Biotechnology) for 30 minutes on ice. Following 3 washes, the cells were resuspended in staining buffer containing Propidium Iodide (PI), a vital stain that allows for the identification of permeable cells, and analyzed by FACS.

5 From these experiments, the results of which are illustrated in Figures 20-21, O8E expression was detected on the surface of transfected HEK293 cells and SKBR3 cells by FACS analysis using rabbit anti-O8E sera. Expression was also detected in transfected HEK293 cell lysates by Western blot analysis (Figure 22).

10

Example 8

Generation and Characterization of anti-O8E mAbs.

11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180

added and the plates incubated for 60 minutes at room temperature. The plates were washed 5 times with washing buffer and the secondary antibody, donkey-anti mouse Ig linked to horseradish peroxidase (HRP)(Jackson ImmunoResearch), was added for 60 minutes. The plates were again washed 5 times in washing buffer, followed by the addition
 5 of the peroxidase substrate. Of the hybridoma clones generated, 15 secreted mAbs that recognized the entire O8E protein. Epitope mapping revealed that of these 15 clones, 14 secreted mAbs that recognized the O8E amino acid residues 61-80 and one clone secreted a mAb that recognized amino acid residues 151-170.

10 For flow cytometric analysis, HEK293 cells which had been stably transfected with O8E and SKBR3 cells which express O8E mRNA, were harvested and washed in flow staining buffer (PBS+1%BSA+Azide). The cells were incubated with the supernatant from the mAb hybrids for 30 minutes on ice followed by 3 washes with staining buffer. The cells were incubated with goat-anti mouse Ig-FITC for 30 minutes on
 15 ice, followed by three washes with staining buffer before being resuspended in wash buffer containing propidium iodide. Flow cytometric analysis revealed that 15/15 mAbs were able to detect O8E protein expressed on the surface of O8E-transfected HEK293 cells. 6/6 mAbs tested on SKBR3 cells were able to recognize surface expressed O8E.

20 Example 9

Extended DNA and protein sequence analysis of sequence O772P

A full-length sequence encompassing clones 3f, 6b, 8e, and 12 was obtained by screening an ovarian tumor (SCID-derived) cDNA library described in detail in
 25 Example 2. This 2996 base pair sequence, designated O772P, is presented in SEQ ID NO: 311, and the encoded 914 amino acid protein sequence is shown in SEQ ID NO: 312. The DNA sequence O772P was searched against public databases including Genbank and showed a significant hit to Genbank Accession number AK024365 (SEQ ID NO: 457). This Genbank sequence was found to be 3557 base pairs in length and encodes a protein

1156 amino acids in length (SEQ ID NO: 459). A truncated version of this sequence, residues 25-3471, in which residue 25 corresponds to the first ATG initiation codon in the *Genbank* sequence, (SEQ ID NO: 456), encodes a protein that is 1148 amino acids in length (SEQ ID NO: 458). The published DNA sequence (SEQ ID NO: 457) differs from
 5 O772P in that it has a 5 base pair insertion corresponding to bases 958-962 of SEQ ID NO: 457. This insertion results in a frame shift such that SEQ ID NO: 457 encodes an additional N-terminal protein sequence relative to O772P (SEQ ID NO: 312). In addition, O772P encodes a unique N-terminal portion contained in residues 1-79 (SEQ ID NO: 460). The N-terminal portion of SEQ ID NO: 456, residues 1-313, also contains unique sequence
 10 and is listed as SEQ ID NO: 461.

Example 10

The Generation of Polyclonal Antibodies for Immunohistochemistry and Flow Cytometric Analysis of the Cell Associated Expression Pattern of Molecule O772P

15 The O772P molecule was identified in Examples 2 and 9 of this application. To evaluate the subcellular localization and specificity of antigen expression in various tissues, polyclonal antibodies were generated against O772P. To produce these antibodies, O772P-1 (amino acids 44-772 of SEQ ID NO:312) and O772P-2 (477-914 of SEQ ID
 20 NO:312) were expressed in an *E. coli* recombinant expression system and grown overnight at 37°C in LB Broth. The following day, 10ml of the overnight culture was added to 500ml of 2xYT containing the appropriate antibiotics. When the optical density of the cultures (560 nanometers) reached 0.4-0.6 the cells were induced with IPTG. Following induction, the cells were harvested, washed, lysed and run through a French Press at a pressure of
 25 16000 psi. The cells were then centrifuged and the pellet checked by SDS-PAGE for the partitioning of the recombinant protein. For proteins that localize to the cell pellet, the pellet was resuspended in 10mM Tris, pH 8.0, 1% CHAPS and the inclusion body pellet washed and centrifuged. The washed inclusion body was solubilized with either 8M urea or 6M guanidine HCL containing 10mM Tris, pH 8.0, plus 10mM imidazole. The

solubilized protein was then added to 5ml of nickel-chelate resin (Qiagen) and incubated for 45 minutes at room temperature.

Following the incubation, the resin and protein mixture was poured through a column and the flow through collected. The column was washed with 10-20 column
 5 volumes of buffer and the antigen eluted using 8M urea, 10mM Tris, pH 8.0, and 300 mM imidazole and collected in 3ml fractions. SDS-PAGE was run to determine which fractions to pool for further purification. As a final purification step, a strong anion exchange resin was equilibrated with the appropriate buffer and the pooled fractions were loaded onto the column. Each antigen was eluted from the column with an increasing salt gradient.
 10 Fractions were collected and analyzed by a SDS-PAGE to determine which fractions from the column to pool. The pooled fractions were dialyzed against 10mM Tris, pH 8.0, and the resulting protein was submitted for quality control for final release. The release criteria were: (a) purity as determined by SDS-PAGE or HPLC, (b) concentration as determined by Lowry assay or Amino Acid Analysis, (c) identity as determined by amino terminal protein,
 15 and (d) endotoxin levels as determined by the Limulus (LAL) assay. The proteins were then filtered through a 0.22µM filter and frozen until needed for immunizations.

To generate polyclonal antisera, 400µg of O772P-1 or O772P-2 was combined with 100µg of muramyl dipeptide (MDP). The rabbits were immunized every 4 weeks with 100µg of antigen mixed with an equal volume of Incomplete Freund's
 20 Adjuvant (IFA). Seven days following each boost, the animals were bled and sera was generated by incubating the blood at 4°C for 12-24 hours followed by centrifugation.

To characterize the antisera, 96 well plates were coated with antigen followed by blocking with BSA. Rabbit sera was diluted in PBS and added to each well. The plates were then washed, and goat anti-rabbit horseradish peroxidase (HRP). The
 25 plates were again washed and TMB microwell Peroxidase Substrate was added. Following this incubation, the colorimetric reaction was stopped and the plates read immediately at 450nm. All polyclonal antibodies showed immunoreactivity to the appropriate antigen.

Immunohistochemistry analysis of O772P expression was performed on paraffin-embedded formalin fixed tissue. O772P was found to be expressed in normal

ovary and ovarian tumor, but not in normal heart, kidney, colon, lung or liver. Additionally, immunohistochemistry and flow cytometric analysis indicates that O772P is a plasma membrane-associated molecule. O772P contains 1 plasma transmembrane domain predicted to be encoded by amino acids 859-880. The N-terminus of O772P is extracellular and is encoded by amino acids 1-859, while the C-terminus is intracellular. Sequence analysis shows that there are 17 potential N-linked glycosylation sites.

Example 11

O772P is Expressed on the Surface of Primary Ovarian Tumor Cells

10

For recombinant expression in mammalian cells, the O772P-21008 (SEQ ID NO:387) and O772P full length cDNA (SEQ ID NO:311 encoding the protein of SEQ ID NO:312) were subcloned into mammalian expression vectors pBIB or pCEP4 respectively. These constructs were transfected into HEK293 cells using Fugene 6 (Roche). The HEK cells were then plated at a density of 100,000 cells/ml in DMEM containing fetal bovine serum (FBS) and grown overnight. The following day, 2 μ l of Fugene 6 was added to 100 μ l of DMEM, which contained no FBS, and incubated for 15 minutes at room temperature. The Fugene 6/DMEM mixture was then added to 1 μ g of O772P/pBIB or O772P/pCEP4 plasmid DNA and incubated for an additional 15 minutes at room temperature. The Fugene 6/DNA mix was then added to the HEK293 cells and incubated for 48-72 hours at 37°C with 7% CO₂. The cells were rinsed and pelleted by centrifugation.

For Western Blot analysis, whole cell lysates were generated by incubating the cells in lysis buffer followed by clarification by centrifugation. The samples were diluted and run on SDS-PAGE. The gel was then transferred to nitrocellulose and probed using purified anti-O772P-2 rabbit polyclonal antibody. The blot was revealed with a goat anti-rabbit Ig coupled to HRP followed by incubation in ECL substrate. Western Blot analysis revealed that O772P-21008 could be detected in HEK293 cells that had been transfected with O772P.

To determine the cell expression profile of O772P in cells, primary ovarian tumor cells were grown in SCID mice. The cells were retrieved from the mice and analyzed by flow cytometry. Briefly, cells washed in cold staining buffer containing PBS, 1% BSA, and Na Azide. The cells were incubated for 30 minutes with 10µg/ml of purified anti-O772P-1 and O772P-2 polyclonal sera. Following this incubation, the cells were washed three times in staining buffer and incubated with goat anti-rabbit Ig (H+L) conjugated to FITC (Southern Biotechnology). The cells were washed and resuspended in staining buffer containing Propidium Iodide (PI), a vital stain that identifies non-viable cells. The cells were then analyzed using Fluorescence Activated Cell Sorting (FACS). FACS analysis revealed that O772P was present on the cells surface. Surface expression of O772P on tumor cells allows for immune targeting by therapeutic antibodies.

Example 12

Functional Characterization of Anti-O8E Monoclonal Antibodies

Mouse monoclonal antibodies (mAb) raised against E. coli derived O8E, as described in Example 8, were tested for their ability to promote O8E antigen internalization. Internalization of the antibody was determined using an in vitro cytotoxicity assay. Briefly, HEK293 and O8E/HEK transfected cells were plated into 96 well plates containing DME plus 10% heat-inactivated FBS in the presence of 50ng/well of purified anti-O8E or control antibodies. The isotype of the anti-O8E mAbs are as follows: 11A6-IgG1/kappa, 15C6-IgG2b/kappa, 18A8-IgG2b/kappa, and 14F1-IgG2a/kappa. W6/32 is a pan anti-human MHC class I mouse monoclonal antibody that serves as a positive control, and two irrelevant mAbs, Ir-Pharm and Ir-Crxa were included as negative controls. Following incubation with the O8E specific antibodies or the relevant controls antibodies, the mAb-zap, a goat anti-mouse Ig-saporin conjugated secondary antibody (Advanced Targeting Systems) was added at a concentration of 100ng/ml to half of the wells, and the plates were incubated for 48 to 72 hours at 37°C in a 7% CO₂ incubator. This assay takes advantage of the toxic nature of saporin, a ribozyme inactivating protein,

which when internalized has a cytotoxic effect. Following incubation with the mAb-zap, internalization was quantitated by the addition of MTS reagent, followed by reading the OD490 of the plate on a microplate ELISA reader. Figure 25 depicts the results from these assays. The top panel represents HEK cells that have not been transfected with O8E and therefore O8E antibody should not bind and be internalized. Levels of proliferation were the same in all samples whether they were incubated with or without the mAb-zap, with the exception of the positive control Ab, W6/32. The lower panel represents cells that have been transfected with O8E and therefore should bind O8E specific antibodies. Antibodies from the hybridomas 11H6, 14F1, and 15C6, which recognize the amino acids 61-80 of O8E were able to promote internalization of the O8E surface protein as measured by decreased levels of proliferation due to the toxic nature of the mAb-zap (See Figure 25). The antibody generated by the hybridoma 18A8, which recognizes amino acids 151-170 of O8E, was unable to promote internalization as determined by normal levels of proliferation either in the absence or presence of the mAb-zap.

15

From the foregoing it will be appreciated that, although specific embodiments of the invention have been described herein for purposes of illustration, various modifications may be made without deviating from the spirit and scope of the invention. Accordingly, the invention is not limited except as by the appended claims.